


1284315



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

February 27, 2006

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/558,504

FILING DATE: March 31, 2004

RELATED PCT APPLICATION NUMBER: PCT/US04/32599

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS US60/558,504




Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

Abstract of the Disclosure

The subject invention concerns polynucleotides encoding a plant 2-phenylethanol dehydrogenase enzyme. In one embodiment, the polynucleotide encodes a tomato 2-phenylethanol dehydrogenase. The subject invention also concerns polynucleotides encoding a
5 plant phenylalanine decarboxylase enzyme. In one embodiment, the polynucleotide encodes a tomato phenylalanine decarboxylase. The subject invention also concerns 2-phenylethanol dehydrogenase polypeptides and phenylalanine decarboxylase polypeptides encoded by polynucleotides of the present invention. The subject invention also concerns methods for providing a plant with an increased flavor and aroma volatile. Plants can be transformed with
10 one or more polynucleotide of the present invention. The subject invention also concerns these transformed plant cells, plant tissue, and plants and transgenic progeny thereof.






FIG. 2

1 GCCCTTCTAA TAGACTCAC TATAGGGCAA GCAGTGGTAA CAACGCAGAG
51 TACGCGGGG AAGGATAATC TCTCAAATTA CTTTCTTTTT TTTTCCATC
101 AATTCCTTAT ACCAAAATAA TATTATTGTT TTTTCTCCT CTGTTCTGCG
151 TTCGTAATTT TGCTGAGAGA AATGAGTGTG ACAGCGAAAA CAGTGTGTGT
201 AACAGGAGCT TCAGGTACA TAGCTTCATG GCTAGTCAAA TTCTTGCTAC
251 ATAGTGGTTA CAATGTGAAG GCTTCTGTTT GTGATCCAAA TGATCCCAAG
301 AAAACGCAGC ACTTGCTTTC TCTTGGTGGG GCCAAGGAGA GGCTTCACCT
351 GTTCAAAGCA AACCTATTAG AAGAAGGTTT ATTTGATGCT GTAGTTGATG
401 GATGTGAAGG TGTATTCCAT ACAGGCTCTC CTTTTTACTA CTCTGTTACA
451 GACCCACAGG CTGAATTACT TGATCCTGCT GTTAAGGGAA CACTCAATCT
501 TCTCGGGTCA TGTGCCAAAG CACCATCAGT AAAACGAGTT GTTTTAACGT
551 CTTCCATAGC TGCAGTTGCT TACAGTGGTC AGCCTCGGAC ACCTGAGGTT
601 GTGGTTGATG AGAGCTGGTG GACCAGTCCA GACTACTGCA AAGAAAAACA
651 GCTCTGGTAT GTCCTCTCAA AGACATTGGC TGAGGATGCT GCGTGGAAAGT
701 TTGTGAAGGA GAAAGGCATT GATATGGTTG TAGTAAACCC TGCTATGGTT
751 ATTGGTCCCTC TGTTACAGCC TACACTTAAT ACCAGTTCTG CTGCAGTCTT
801 GAGCTTGGTA AATGGTGTG AGACATACCC AAATTCCTCT TTTGGGTGGG
851 TTAACGTGAA AGATGTTGCA AATGCACATA TTCTTGCATT TGAGAACCCCT
901 TCAGCTAATG GGAGATACTT AATGGTTGAG AGGGTTGCAC ACTATTCTGA
951 TATATTGAAG ATATTGCGTG ACCTTTATCC TACTATGCAA CTTCCAGAAA
1001 AGTGTGCTGA TGACAACCCA TTGATGCAAA ATTATCAAGT ATCAAAGGAG
1051 AAGGCAAAA GCTTGGGTAT TGAGTTTACT ACCCTTGAAG AAAGCATCAA
1101 AGAACTGTT GAAAGTTTGA AGGAAAAGAA GTTTTTTGGG GGTTCATCTT
1151 CTATGTAAAA GGCTTCTCAA AGCTTTTATG GTTTTGTGA ACAATACTAC
1201 CCACCCACC CTACCCCTACA CACTTTTTTT TTTTACTTCT TTTAGCTAAT
1251 TATAGAAATCA AGAAGTCGAA TGGTATATCC GTTAATAAAT TTCGATCAGA
1301 TGAGGTTGAA ATTTGTTCTA TATCTAGAGA TTTTACAGA CTGGTTTGAT
1351 AGAAAAAAA AAAAAA

FIG. 3A

1 MSVTAKTVCV TGASGYIASW LVKELLHSGY NVKASVRDPN DPKKTQHLLS
51 LGGAKERLHL FKANLLEEGS FDAVVDGCEG VFHTASPFYY SVTDPQAE LL
101 DPAVKGTLNL LGSCAKAPSV KRVLTSIA AVAYSGQPT PEVVDES WW
151 TSPDYCKEQ LWVLSKTIA EDAAWKFVKE KGIDMVVNP AMVIGPLLQP
201 TLNTSSAAVL SLVNGAETYP NSSFGWNVK DVANAHILAF ENPSANGRYL
251 MVERVAHYSD ILKILRDLYP TMQLPEKCAD DNPLMQNYQV SKEKAKSLGI
301 EFTTLEESIK ETVESLKEKK FFGSSSM

FIG. 3B




FIG. 4




FIG. 5A-1

Phenylacetaldehyde




FIG. 5A




FIG. 5B-1

Cinnamaldehyde




FIG. 5B




FIG. 5C-1

Vanillin




FIG. 5C




FIG. 6A




FIG. 6B




FIG. 7

1 ATGGGAAGTT TATCATTGA GAAGGATTTT GAGCCATCAG CAATTACTCC
51 AAGAGGATTA GCACCACCTG GATTAATGT AAATGGTGAT TTTGGTGA
101 TGATGAGACT TAAGGTGCA TCAACACCAA CAACACCAAG AAAAACTTG
151 AATCTTTCAG TGACGGAGCC AGGAAAAAAT GATGGACCTA GTTTGGATTG
201 TACATTGATG AATTATATTG ATACACTCAC CCAACGTATC AACTATCATA
251 TCGGTTATCC AGTTAACATA TGTATGAGC ACTATGCTAA TTTAGCCCCA
301 CTTTTACAAT TTCATTTAAA TAATTGTGTT GATCCATTTC TTCAAAAATAC
351 TGTGGATTTT CATTCAAAGG ATTTTGAAGT GGCTGTTTTA AATTGGTTTG
401 CTGATTTATG GGAATTGAA AGAGATCAAT ATTGGGGCTA TGTAACAAAT
451 GGTGGTACTG AAGGAAATTT ACATGGCATT TTGGTTGGGA GAGAAATTGTT
501 TCCAGATGGA ATTTTATATG CATCAAAAGA CTCTCATTAC TCAGTGGCTA
551 AGGCAGCAAT GATGTATAGA ATGGATTTTG AAAATATTAA CGCATCAATA
601 AATGGAGAAA TCGATTATTC TGATTTGAAA GTTAAATTAC TTCAAAAACAA
651 GGGAAAACCA CGGATAATTA ATGTTACAAT TGGCACTACT TTAAAGGAG
701 CTGTTGATGA TCTTGATGTT ATTCCTCAA TACTTGAAGA GTGTGGTTAC
751 ACACGAGATC AATTTTATAT TCATTGTGAT GCAGCACTAA ATGGACTTAT
801 TATTCTTTT ATTAAAAATA TGATTACTTT CAAGAAGCCA ATTGGAAGTG
851 TGACAAATTC TGGTCACAAG TTTTGGGAT GTCCAATGCC TTGTGGAGTT
901 CAAATAACAA GGAAAAGTTA CATTAAATAC CTTTCGAGAA GAGTCGAATA
951 TATTGCTTCT GTGGATGCTA CAATTTCTGG AAGTCGAAAT GGTTTGACTC
1001 CGATCTTCTT ATGGTACAGT ATAAGTGCTA AAGTCAAAT TGGTTTTCAG
1051 AAAGACGTTA AGAGATGTTT TGACAATGCT AAGTACTTGA AAGACCGTCT
1101 TCAGCAAGCA GGAATCAGCG TCATGCTGAA TGAGCTTAGC ATCATAGTTG
1151 TCCTCGAGAG GCCTCGTGAC CATGAATTCG TTCGTCGTTg GCAATTATCT
1201 TGTGTGAGAG ATAATGCACA TGTATTGTT ATGCCAGGCA TAAC TAGAGA
1251 AACTCTTGAT GGTTTTATTA ATGATTGCT TCAACAAAGG AAAAAATGGT
1301 ATCAAGATGG AAGAATTAGC CCTCCTTG TGCAAAATGA TATTGGTGCT
1351 CAAAATTGTG CTTGCTCTTA TCATAAAAT GATTACATTA TTGCTTAG

FIG. 8A

1	M	G	S	L	S	F	E	K	D	F	E	P	S	A	I	T	P	R	G	L	A	P	P	G	L	I	V	N	G	D	F	G	E	M	R	L	K	V	S	S	T	P	T	T	P	R	K	N	L		
51	N	L	S	V	T	E	P	G	K	N	D	G	P	S	L	D	C	T	I	M	N	Y	I	D	T	I	L	T	Q	R	I	N	Y	H	I	G	Y	P	V	N	I	C	Y	E	H	Y	A	N	L	A	P
101	L	L	Q	F	H	L	N	N	C	G	D	P	F	L	Q	N	T	V	D	F	H	S	K	D	F	E	V	A	V	L	N	W	F	A	D	L	W	E	I	E	R	D	Q	Y	W	G	Y	V	T	N	
151	G	G	T	E	G	N	L	H	G	I	L	V	G	R	E	L	F	P	D	G	I	L	Y	A	S	K	D	S	H	Y	S	V	A	K	A	A	M	M	Y	R	M	D	F	E	N	I	N	A	S	I	
201	N	G	E	I	D	Y	S	D	L	K	V	K	L	L	Q	N	K	G	K	P	A	I	N	V	T	I	G	T	T	F	K	G	A	V	D	D	L	D	V	I	L	Q	I	L	E	E	C	G	Y		
251	T	R	D	Q	F	Y	I	H	C	D	A	A	L	N	G	L	I	I	P	F	I	K	N	M	I	T	F	K	K	P	I	G	S	V	T	I	S	G	H	K	F	L	G	C	P	M	P	C	G	V	
301	Q	I	T	R	K	S	Y	I	N	N	L	S	R	R	V	E	Y	I	A	S	V	D	A	T	I	S	G	S	R	N	G	L	T	P	I	F	L	W	Y	S	I	S	A	K	G	I	G	F	Q		
351	K	D	V	K	R	C	F	D	N	A	K	Y	L	K	O	R	L	Q	Q	A	G	I	S	V	M	L	N	E	L	S	I	I	V	V	L	E	R	P	R	D	H	E	F	V	R	R	W	Q	L	S	
401	C	V	R	D	M	A	H	V	I	V	M	P	G	I	T	R	E	T	L	D	G	F	I	N	D	L	L	Q	Q	R	K	K	W	Y	Q	D	G	R	I	S	P	P	C	V	A	N	D	I	G	A	
451	Q	N	C	A	C	S	Y	H	K	I	D	Y	I	I	A																																				

FIG. 8B

1 ATGGGTAGTC TCTCACTTGA AATGGATTTT GAGCCATCAC CCATGACACC
51 CAGAAAGTTTA GCAGCGATGA CACCTAGAAG TTTAGCGCGA CGACGATTGT
101 TTCCGAACGT GGACAACAAG AAACAGAAAA TGGCACAACC AGGTGCAGGA
151 CCAAGGAAGA ACTTGGAAC TGAAGTCATG GAGCCTGCGT TGAAGAATGA
201 TGGTCCCTTCT TTGGACACTA TCTTGGTTAA TTATTTGGAC ACACTTACAC
251 AACGAGTCAA TTATCATTTA GGTATATCCAG TCAACATATG TTATGATCAC
301 TATGCAACGC TAGCACCAC TTTGCAGTTT CACCTAAACA ATTGTGGTGA
351 TCCTTTCCCTA CAAAATACTG TCGATTTCCA TTCTAAAGAC TTTGAAAGTGG
401 CTGTTTTGAA TTGGTTTGCA AAACITTTGGG AAATTGAAAA GGATCAATAT
451 TGGGGATATG TTACCAATGG TGGCACCAGG GGCAATCTCC ATGGTATTTT
501 GTTAGGGAGA GAGCTACTTC CTGAAGGAAT ATTATATGCA TCAAAAAGACT
551 CTCATTACTC AGTATTCAAA GCTGCAAGAA TGTATAGAAT GGATTCAGAA
601 ACAATCAACA CATCAGTAAA TGGAGAGATG GATTATTCAG ATTTAAGAGC
651 AAAGTTACTT CAAAATAAGG ATAAACCAGC TATTATAAAT GTCACAATTG
701 GAACTACATT CAAAGGAGCA ATCGATGACC TGGATGTTAT TCTTGAAATA
751 CTCAAAAGAAT GTGGCTATTC ACAAGATCGA TTTTACATTC ACTGTGATGC
801 AGCACTATGT GGTCTTATGA CCCCTTTTAT AAACAATATG ATTAGTTTCA
851 AGAAGCCAAT TGGAAAGTGC ACAATTTCGT GACACAAGTT TTTGGGATGT
901 CCAATGCCTT GTGGTGTCCA AATAACAAGA AAAAGCTACA TCAATAATCT
951 CTCACACAAT GTGGAATACA TTGCTTCTGT GGAIGCCACT ATTTCTGGTA
1001 GCCGTAACGG TTAACTCCA ATTTTCTTAT GGTATAGCTT GAGCGCAAAA
1051 GGTCAGTTG GACTTCAAAA GGATGTTAAA AGATGCTCG ACAATGCCAA
1101 ATATTTGAAA GATCGTCTTC AACAAAGCAGG GATAAGTGC ATGCTGAATG
1151 AGCTAAGCAT CATAGTTGTA CTTGAAAGGC CTCGTGACCA TGAATTTGTG
1201 CGTCGTTGGC AACTCTCATG CGTCAAGGAT ATGGCACATG TTATTGTGAT
1251 GCCAGGAATC ACACGAGAAA TGCTTGACAA CTTCATGAGT GAATTAGTGC
1301 AACAAAGAAA AGTATGGTAT CAAAATGGAA AGACTGATCC TCCTTGTGTT
1351 GGAGAGGATA TTGGTGCTCA AAATTGTGCA TGCTCTTATC ATAAGATTGA
1401 CTACATCTGT CCTTAG

FIG. 9A

1 MGSLSLEMDF EPSPMTPRSL AAMTPRSLAR RRLFPNVNDK KQKMAQPGAG
51 PRKNLELEVMEPALKNDGPS LDTILVNYLD TLTQRVNYHL GYPVNICYDH
101 YATLAPLLQF HLNCGDPFL QNTVDFHSKD FEVAVLNWFA KLWEIEKDQY
151 WGYVTNGGTE GNLHGILLGR ELLPEGILYA SKDSHYSVFK AARMYRMDSE
201 TINTSVNGEM DYSDLRAKLL QNKDKPAIIN VTIGTTFKGA IDDLDVILEI
251 LKECGYSQDR FYIHCDAAALC GLMTPFINNM ISFKKPIGSV TISGHKFLGC
301 PMPCGVQITR KSYINNLSN VEYIASVDAT ISGSRNGLTP IFLWYLSAK
351 GQVGLQKDVK RCLDNAKYLK DRLQQAGISV MLNELSIIVV LERPRDHEFV
401 RRWQLSCVKD MAHVIVMPGI TREMLDNFMS ELVQQQRKVWY QNGKTDPPCV
451 GEDIGAQNCA CSYHKIDYIC P

FIG. 9B




FIG. 10

1 ATGGGTAGTC TCTCACTTGA AATGGATTTT GAGCCATCAC CTATGACACC
51 CAGAAGTTTA GCAGCGATGA CACCTAGAAG TTTAGCGGG CGAAGATTGT
101 TTCCCAATGT GGACAACAAA AAACAAAAGG TGCAACAATC AGGTGCAGGG
151 CCAAGGAAGA ACTTACAAC TGAAGTCATG GAACCTGCAT TGAACAATGC
201 TGGTCCCTCT TTGGACACTA TATTGGTCAA TTATTTAGAC ACACTTACAC
251 AACGAGTCAA TTATCATTTA GGTATCCAG TCAACATTG TTATGATCAC
301 TATGCAACTT TAGCACCAC TTTACAGTTT CACCTAAACA ATTGTGGTGA
351 TCCTTTCCTA CAAAACACTG TCGATTTCCA TTCTAAAGAC TTTGAAGTGG
401 CTGTTTTGAA TTGGTTTGCA AAACATATGG AAATTGAAAA GGATCAATAC
451 TGGGGATATG TTACCAATGG TGGCACCGAA GGCAATCTCC ATGGTATTTT
501 GTTAGGGAGA GAGCTACTTC CTGATGGAAT ATTATATGCG TCAAAAAGACT
551 CTCACATATC GGTCTTCAAA GCTGCAAGAA TGTATAGAAT GGATTCAGAA
601 ACAATCAACA CATCAGTAAA CGGAGAGATG GATTATTAG ATTAAAGAGC
651 AAAGTTACTT CAAAATAAGG ATAAACCAGC TATTATAAAT GTCACAATTG
701 GAACTACGTT CAAAGGAGCA ATCGATGACC TGGATGTTAT TCTTGAAAAA
751 CTCAAAAGAA GTGGCTATTC GCAAGATAGG TTTTACATCC ACTGTGATGC
801 TGCACTATGT GGTCTTATGA CCCCTTTTAT AAACAATATG ATTAGTTTCA
851 AGAAGCCAA TGGAAAGTGC ACAATTTCTG GACACAAGTT TTTGGGATGT
901 CCAATGCCCTT GTGGTGTCCA AATTACAAGA AAGAGTTACA TCAATAATCT
951 CTCACAACAA GTGGAATACA TTGCTTCTGT CGATGCCACT ATTTCTGGCA
1001 GCCGTAACGG TTTAACTCCA ATTTTCTTGT GGTATAGCTT GAGCGCAAAA
1051 GGTCAAGTTG GACTTCAAAA GGATGTTAAA AGATGTCTCG ACAATGCCAA
1101 ATATTTGAAA GATCGTCTTC AAAAAGCAGG AATAAGTGC ATGTTAAATG
1151 AGCTTAGCAT CATAGTTGTA CTTGAAAGGC CTCGTGACCA TGAATTTGTC
1201 CGTCGTTGGC AACTCTCATG CGTCAAGGAT ATGGCACATG TTATTGTAAT
1251 GCCAGGCATC ACACGAGAAA TGCTTGACAA TTTACAGAGT GAATTAGTGC
1301 AACAAAGAAA AGTATGGTAT CAAAATGGAC AGACCAATCC TCCTTGTGTT
1351 GGAGAGGATA TTGGTGCTCA AAATTGTGCA TGCTCTTATC ATAAGATTGA
1401 CTACATCTGT CCTTAG

FIG. 11A

1	MGSLSEMDF	EPSPMTPRSL	AAMTPRSLAR	RRLFPNVNDNK	KQKVQQSGAG
51	PRKNLQLEVM	EPALNNAGPS	LDTILVNYLD	TLTQRVNYHL	GYPVNICYDH
101	YATLAPLLQF	HLNCGDPFL	QNTVDFHSD	FEVAVLNWFA	KLWEIEKDQY
151	WGYVTNGGTE	GNLHGILLGR	ELLPGDILYA	SKDSHYSVFK	AARMYRMDSE
201	TINTSVNGEM	DYSDLRAKLL	QNKDKPAIIN	VTIGTTFKGA	IDDLDVILET
251	LKECGYSQDR	FYIHCDAAALC	GLMTPFINNM	ISFKKPIGSV	TISGHKFLGC
301	PMPCGVQITR	KSYINNLSIN	VEYIASVDAT	ISGSRNGLTP	IFLWYSLSAK
351	GQVGLQKDVK	RCLDNAKYLK	DRLQKAGISV	MLNELSIIVV	LERPRDHEFV
401	RRWQLSCVKD	MAHVIVMPGI	TREMLDNFTS	ELVQQRKVWY	QNGQTNPPCV
451	GEDIGAQNCA	CSYHKIDYIC	P		

FIG. 11B

1		50
Lp-cLEC73K23	MGSLSLEMDF	EPSPMTPRSL AAMTPRSLAR RRLFPNVDNK KQKVQQSGAG
Le-cLEC73K23	MGSLSLEMDF	EPSPMTPRSL AAMTPRSLAR RRLFPNVDNK KQKMAQPGAG
Le-cLEC75E21	MGSLSFEKDF	EPSAITPRGL A...PPGLIV NGDFGEM..M RLVSSSTPTT
51		100
Lp-cLEC73K23	PRKNLQLEVM	EPALNNAGPS LDTILVNYLD TLTQRVNYHL GYPVNICYDH
Le-cLEC73K23	PRKNLELEVM	EPALKNDGPS LDTILVNYLD TLTQRVNYHL GYPVNICYDH
Le-cLEC75E21	PRKNLNLSVT	EPG.KNDGPS LDCTLMNYID TLTQRINYHI GYPVNICYEH
101		150
Lp-cLEC73K23	YATLAPLLQF	HLNNCGDPFL QNTVDFHSKD FEVAVLNWFA KLWEIEKDQY
Le-cLEC73K23	YATLAPLLQF	HLNNCGDPFL QNTVDFHSKD FEVAVLNWFA KLWEIEKDQY
Le-cLEC75E21	YANLAPLLQF	HLNNCGDPFL QNTVDFHSKD FEVAVLNWFA DLWEIERDQY
151		200
Lp-cLEC73K23	WGYVTNGGTE	GNLHGILLGR ELLPDGILYA SKDSHYSVFK AARMYRMDSE
Le-cLEC73K23	WGYVTNGGTE	GNLHGILLGR ELLPEGILYA SKDSHYSVFK AARMYRMDSE
Le-cLEC75E21	WGYVTNGGTE	GNLHGILVGR ELFPDGILYA SKDSHYSVAK AAMMYRMDFE
201		250
Lp-cLEC73K23	TINTSVNGEM	DYSDLRAKLL QNKDKPAIN VTIGTTFKGA IDDLDVILET
Le-cLEC73K23	TINTSVNGEM	DYSDLRAKLL QNKDKPAIN VTIGTTFKGA IDDLDVILEI
Le-cLEC75E21	NINASINGEI	DYSDLKVLL QNKGKPAIN VTIGTTFKGA VDDLVDVILQI

FIG. 12A

251		300
Lp-cLEC73K23	LKECGYSQDR FYIHCDAAALC GLMTPFINNM ISFKKPIGSV TISGHKFLGC	
Le-cLEC73K23	LKECGYSQDR FYIHCDAAALC GLMTPFINNM ISFKKPIGSV TISGHKFLGC	
Le-cLEC75E21	LEECGYTRDQ FYIHCDAAALN GLIIPFIKIM ITFKKPIGSV TISGHKFLGC	
301		350
Lp-cLEC73K23	PMPCGVQITR KSYINNLSLN VEYIASVDAT ISGSRNGLTP IFLWYSLSAK	
Le-cLEC73K23	PMPCGVQITR KSYINNLSLN VEYIASVDAT ISGSRNGLTP IFLWYSLSAK	
Le-cLEC75E21	PMPCGVQITR KSYINNLSRR VEYIASVDAT ISGSRNGLTP IFLWYSLSAK	
351		400
Lp-cLEC73K23	GQVGLQKDVK RCLDNAKYLK DRLLQAGISV MLNELSIIVV LERPRDHEFV	
Le-cLEC73K23	GQVGLQKDVK RCLDNAKYLK DRLLQAGISV MLNELSIIVV LERPRDHEFV	
Le-cLEC75E21	GQIGFQKDVK RCFDNAKYLK DRLLQAGISV MLNELSIIVV LERPRDHEFV	
401		450
Lp-cLEC73K23	RRWQLSCVKD MAHVIVMPGI TREMLDNFTS ELVQQRKVWY QNGQTNPPCV	
Le-cLEC73K23	RRWQLSCVKD MAHVIVMPGI TREMLDNFMS ELVQQRKVWY QNGKTDPPCV	
Le-cLEC75E21	RRWQLSCVRD MAHVIVMPGI TRETLDDGFN DLLQQRKKWY QDGRISPPCV	
451		472
Lp-cLEC73K23	GEDIGAQNCA CSYHKIDYIC P	
Le-cLEC73K23	GEDIGAQNCA CSYHKIDYIC P	
Le-cLEC75E21	ANDIGAQNCA CSYHKIDYII A	

FIG. 12B






FIG. 13A



Retention time (mins)

FIG. 13B



Retention time (mins)

FIG. 13C

SEQUENCE LISTING

<110> Klee, Harry J.
Tieman, Denise

<120> Isolation of a Gene Involved in Synthesis of a Flavor and Aroma
Volatile in Plants

<130> UF-386CP

<160> 9

<170> PatentIn version 3.2

<210> 1

<211> 1367

<212> DNA

<213> Lycopersicon esculentum

```

<400> 1
gcccttctaa tacgactcac tatagggcaa gcagtggtaa caacgcagag tacgcggggg      60
aaggataatc tctcaaatta ctttcttttt ttttcctatc aattctttat accaaaataa     120
tattattggt tttttctcct ctgtttctgc ttcgtatttt tgctgagaga aatgagtgtg     180
acagcgaaaa cagtgtgtgt aacaggagct tcaggttaca tagcttcatt gctagtcaaa     240
ttcttgctac atagtgggta caatgtgaag gcttctgttc gtgatccaaa tgatcccaag     300
aaaacgcagc acttgctttc tcttggtggg gccaaaggaga ggcttcactt gttcaaagca     360
aacctattag aagaagggtc atttgatgct gtagttgatg gatgtgaagg tgtattccat     420
acagcgtctc ctttttacta ctctgttaca gaccacagg ctgaattact tgatcctgct     480
gttaagggaa cactcaatct tctcgggtca tgtgccaaag caccatcagt aaaacgagtt     540
gttttaacgt cttccatagc tgcagttgct tacagtgggc agcctcggac acctgaggtt     600
gtggttgatg agagctgggt gaccagtcca gactactgca aagaaaaaca gctctggtat     660
gtcctctcaa agacattggc tgaggatgct gcgtggaagt ttgtgaagga gaaaggcatt     720
gatatgggtg tagtaaacc cttctatggt attggtcctc tggtacagcc tacacttaat     780
accagttctg ctgcagtcct gagcttggtg aatggtgctg agacataccc aaattcctct     840
tttggtggg ttaacgtgaa agatggtgca aatgcacata ttcttgcat tgagaaccct     900
tcagctaatt ggagatactt aatggttgag aggggtgcac actattctga tatattgaag     960
atattgcgtg acctttatcc tactatgcaa cttccagaaa agtgtgctga tgacaacca    1020
ttgatgcaaa attatcaagt atcaaaggag aaggcaaaaa gcttggttat tgagtttact    1080

```

```

acccttgaag aaagcatcaa agaaactggt gaaagtttga aggaaaagaa gttttttgga 1140
ggttcatctt ctatgtaaaa ggcttctcaa agcttttatg gttttgttga acaatactac 1200
ccacccccacc ctaccctaca cacttttttt ttttacttct tttagctaata tatagaatca 1260
agaagtcgaa tggatatatcc gttaataaat ttgatcaga tgagggttgaa atttgttcta 1320
tatctagaga tttttacaga ctgggttgat agaaaaaaaa aaaaaaa 1367

```

```

<210> 2
<211> 328
<212> PRT
<213> Lycopersicon esculentum

```

```

<400> 2

```

```

Met Ser Val Thr Ala Lys Thr Val Cys Val Thr Gly Ala Ser Gly Tyr
1          5          10          15
Ile Ala Ser Trp Leu Val Lys Phe Leu Leu His Ser Gly Tyr Asn Val
20          25          30
Lys Ala Ser Val Arg Asp Pro Asn Asp Pro Lys Lys Thr Gln His Leu
35          40          45
Leu Ser Leu Gly Gly Ala Lys Glu Arg Leu His Leu Phe Lys Ala Asn
50          55          60
Leu Leu Glu Glu Gly Ser Phe Asp Ala Val Val Asp Gly Cys Glu Gly
65          70          75          80
Val Phe His Thr Ala Ser Pro Phe Tyr Tyr Ser Val Thr Asp Pro Gln
85          90          95
Ala Glu Leu Leu Asp Pro Ala Val Lys Gly Thr Leu Asn Leu Leu Gly
100         105         110
Ser Cys Ala Lys Ala Pro Ser Val Lys Arg Val Val Leu Thr Ser Ser
115         120         125
Ile Ala Ala Val Ala Tyr Ser Gly Gln Pro Arg Thr Pro Glu Val Val
130         135         140
Val Asp Glu Ser Trp Trp Thr Ser Pro Asp Tyr Cys Lys Glu Lys Gln
145         150         155         160
Leu Trp Tyr Val Leu Ser Lys Thr Leu Ala Glu Asp Ala Ala Trp Lys
165         170         175
Phe Val Lys Glu Lys Gly Ile Asp Met Val Val Val Asn Pro Ala Met
180         185         190

```

Val Ile Gly Pro Leu Leu Gln Pro Thr Leu Asn Thr Ser Ser Ala Ala
 195 200 205

Val Leu Ser Leu Val Asn Gly Ala Glu Thr Tyr Pro Asn Ser Ser Phe
 210 215 220

Gly Trp Val Asn Val Lys Asp Val Ala Asn Ala His Ile Leu Ala Phe
 225 230 235 240

Glu Asn Pro Ser Ala Asn Gly Arg Tyr Leu Met Val Glu Arg Val Ala
 245 250 255

His Tyr Ser Asp Ile Leu Lys Ile Leu Arg Asp Leu Tyr Pro Thr Met
 260 265 270

Gln Leu Pro Glu Lys Cys Ala Asp Asp Asn Pro Leu Met Gln Asn Tyr
 275 280 285

Gln Val Ser Lys Glu Lys Ala Lys Ser Leu Gly Ile Glu Phe Thr Thr
 290 295 300

Leu Glu Glu Ser Ile Lys Glu Thr Val Glu Ser Leu Lys Glu Lys Lys
 305 310 315 320

Phe Phe Gly Gly Ser Ser Ser Met
 325

<210> 3
 <211> 35
 <212> DNA
 <213> Artificial sequence

<220>
 <223> PCR primer

<400> 3
 tccttgcccc caccaagaga aagcaagtgc tgcgt

35

<210> 4
 <211> 1398
 <212> DNA
 <213> Lycopersicon esculentum

<400> 4
 atgggaagtt tatcatttga gaaggatttt gagccatcag caattactcc aagaggatta 60
 gcaccacctg gattaattgt aaatggtgat ttggtgaaa tgatgagact taaggtgtca 120
 tcaacaccaa caacaccaag aaaaaacttg aatctttcag tgacggagcc aggaaaaaat 180
 gatggaccta gtttggtattg tacattgatg aattatattg atacactcac ccaacgtatc 240
 aactatcata tcggttatcc agttaacata tgttatgagc actatgctaa tttagcccca 300

```

cttttacaat ttcatttaaa taattgtggt gatccatttc ttcaaaatac tgtggatttt 360
cattcaaagg attttgaagt ggctgtttta aattggtttg ctgatttatg ggaaattgaa 420
agagatcaat attggggcta tgtaacaaat ggtggtactg aaggaaattt acatggcatt 480
ttggttggga gagaattggt tccagatgga attttatatg catcaaaaga ctctcattac 540
tcagtggcta aggcagcaat gatgtataga atggattttg aaaatattaa cgcacataa 600
aatggagaaa tcgattattc tgatttgaaa gttaaattac ttcaaaacaa gggaaaacca 660
gcgataatta atgttacaat tggcactact tttaaaggag ctgttgatga tcttgatggt 720
attcttcaaa tacttgaaga gtgtggttac acacgagatc aattttatat tcattgtgat 780
gcagcactaa atggacttat tattcctttt attaaaaata tgattacttt caagaagcca 840
attggaagtg tgacaatttc tggtcacaag tttttgggat gtccaatgcc ttgtggagtt 900
caaataacaa ggaaaagtta cattaataac ctttcgagaa gagtcgaata tattgcttct 960
gtggatgcta caatttctgg aagtcgaaat ggtttgactc cgatcttctt atggtacagt 1020
ataagtgcta aaggtcaaat tggttttcag aaagacgta agagatgttt tgacaatgct 1080
aagtacttga aagaccgtct tcagcaagca ggaatcagcg tcatgctgaa tgagcttagc 1140
atcatagttg tctcagagag gcctcgtgac catgaattcg ttcgtcgttg gcaattatct 1200
tgtgtgagag atatggcaca tgttattggt atgccaggca taactagaga aactcttgat 1260
ggttttatta atgatttgct tcaacaaagg aaaaaatggt atcaagatgg aagaattagc 1320
cctccttggt ttgcaaatga tattggtgct caaaattgtg cttgctctta tcataaaatt 1380
gattacatta ttgcttag 1398

```

```

<210> 5
<211> 465
<212> PRT
<213> Lycopersicon esculentum

```

```

<400> 5

```

```

Met Gly Ser Leu Ser Phe Glu Lys Asp Phe Glu Pro Ser Ala Ile Thr
1           5           10           15

```

```

Pro Arg Gly Leu Ala Pro Pro Gly Leu Ile Val Asn Gly Asp Phe Gly
20           25           30

```

```

Glu Met Met Arg Leu Lys Val Ser Ser Thr Pro Thr Thr Pro Arg Lys
35           40           45

```

```

Asn Leu Asn Leu Ser Val Thr Glu Pro Gly Lys Asn Asp Gly Pro Ser
 50                               55                               60

Leu Asp Cys Thr Leu Met Asn Tyr Ile Asp Thr Leu Thr Gln Arg Ile
65                               70                               75                               80

Asn Tyr His Ile Gly Tyr Pro Val Asn Ile Cys Tyr Glu His Tyr Ala
                               85                               90                               95

Asn Leu Ala Pro Leu Leu Gln Phe His Leu Asn Asn Cys Gly Asp Pro
                               100                               105                               110

Phe Leu Gln Asn Thr Val Asp Phe His Ser Lys Asp Phe Glu Val Ala
                               115                               120                               125

Val Leu Asn Trp Phe Ala Asp Leu Trp Glu Ile Glu Arg Asp Gln Tyr
                               130                               135                               140

Trp Gly Tyr Val Thr Asn Gly Gly Thr Glu Gly Asn Leu His Gly Ile
145                               150                               155                               160

Leu Val Gly Arg Glu Leu Phe Pro Asp Gly Ile Leu Tyr Ala Ser Lys
                               165                               170                               175

Asp Ser His Tyr Ser Val Ala Lys Ala Ala Met Met Tyr Arg Met Asp
                               180                               185                               190

Phe Glu Asn Ile Asn Ala Ser Ile Asn Gly Glu Ile Asp Tyr Ser Asp
                               195                               200                               205

Leu Lys Val Lys Leu Leu Gln Asn Lys Gly Lys Pro Ala Ile Ile Asn
                               210                               215                               220

Val Thr Ile Gly Thr Thr Phe Lys Gly Ala Val Asp Asp Leu Asp Val
225                               230                               235                               240

Ile Leu Gln Ile Leu Glu Glu Cys Gly Tyr Thr Arg Asp Gln Phe Tyr
                               245                               250                               255

Ile His Cys Asp Ala Ala Leu Asn Gly Leu Ile Ile Pro Phe Ile Lys
                               260                               265                               270

Asn Met Ile Thr Phe Lys Lys Pro Ile Gly Ser Val Thr Ile Ser Gly
                               275                               280                               285

His Lys Phe Leu Gly Cys Pro Met Pro Cys Gly Val Gln Ile Thr Arg
                               290                               295                               300

Lys Ser Tyr Ile Asn Asn Leu Ser Arg Arg Val Glu Tyr Ile Ala Ser
305                               310                               315                               320

Val Asp Ala Thr Ile Ser Gly Ser Arg Asn Gly Leu Thr Pro Ile Phe
                               325                               330                               335

```

Leu Trp Tyr Ser Ile Ser Ala Lys Gly Gln Ile Gly Phe Gln Lys Asp
 340 345 350

Val Lys Arg Cys Phe Asp Asn Ala Lys Tyr Leu Lys Asp Arg Leu Gln
 355 360 365

Gln Ala Gly Ile Ser Val Met Leu Asn Glu Leu Ser Ile Ile Val Val
 370 375 380

Leu Glu Arg Pro Arg Asp His Glu Phe Val Arg Arg Trp Gln Leu Ser
 385 390 395 400

Cys Val Arg Asp Met Ala His Val Ile Val Met Pro Gly Ile Thr Arg
 405 410 415

Glu Thr Leu Asp Gly Phe Ile Asn Asp Leu Leu Gln Gln Arg Lys Lys
 420 425 430

Trp Tyr Gln Asp Gly Arg Ile Ser Pro Pro Cys Val Ala Asn Asp Ile
 435 440 445

Gly Ala Gln Asn Cys Ala Cys Ser Tyr His Lys Ile Asp Tyr Ile Ile
 450 455 460

Ala
 465

<210> 6
 <211> 1416
 <212> DNA
 <213> Lycopersicon esculentum

<400> 6
 atgggtagtc tctcacttga aatggatttt gagccatcac ccatgacacc cagaagttaa 60
 gcagcgatga cacctagaag tttagcgcgga cgacgattgt ttccgaacgt ggacaacaag 120
 aaacagaaaa tggcacaacc aggtgcagga ccaaggaaga acttggaact tgagggtcatg 180
 gagcctgcat tgaagaatga tggtccttct ttggacacta tcttggttaa ttatttggac 240
 acatttacac aacgagtcaa ttatcattta ggttatccag tcaacatatg ttatgatcac 300
 tatgcaacgc tagcaccact tttgcagttt cacctaaaca attgtggtga tcctttccta 360
 caaaatactg tcgatttcca ttctaaagac tttgaagtgg ctgttttgaa ttggtttgca 420
 aaactttggg aaattgaaaa ggatcaatat tggggatatg ttaccaatgg tggcaccgaa 480
 ggcaatctcc atggtatttt gttagggaga gagctacttc ctgaaggaat attatatgca 540
 tcaaaagact ctctattactc agtattcaaa gctgcaagaa tgtatagaat ggattcagaa 600
 acaatcaaca catcagtaaa tggagagatg gattattcag atttaagagc aaagttactt 660

```

caaaataagg ataaaccagc tattataaat gtcacaattg gaactacatt caaaggagca 720
atcgatgacc tggatgttat tcttgaaata ctcaaagaat gtggctattc acaagatcga 780
ttttacattc actgtgatgc agcactatgt ggtcttatga ccccttttat aaacaatatg 840
attagtttca agaagccaat tggaagtgtc acaatttctg gacacaagtt tttgggatgt 900
ccaatgcctt gtggtgtcca aataacaaga aaaagctaca tcaataatct ctcaacaaat 960
gtggaataca ttgcttctgt ggatgccact atttctggta gccgtaacgg ttttaactcca 1020
attttcttat ggtatagctt gagcgcaaaa ggtcaagttg gacttcacaaa ggatgttaaa 1080
agatgtctcg acaatgccaa atatttgaaa gatcgtcttc aacaagcagg gataagtgtc 1140
atgctgaatg agctaagcat catagttgta cttgaaaggc ctcgtgacca tgaatttgtg 1200
cgtcgttggc aactctcatg cgtcaaggat atggcacatg ttattgtgat gccaggaatc 1260
acacgagaaa tgcttgacaa cttcatgagt gaattagtgc aacaaagaaa agtatggtat 1320
caaatggaa agactgatcc tccttggtgt ggagaggata ttggtgctca aaattgtgca 1380
tgctcttatc ataagattga ctacatctgt ccttag 1416

```

```

<210> 7
<211> 471
<212> PRT
<213> Lycopersicon esculentum

```

```

<400> 7

```

```

Met Gly Ser Leu Ser Leu Glu Met Asp Phe Glu Pro Ser Pro Met Thr
1           5           10          15

Pro Arg Ser Leu Ala Ala Met Thr Pro Arg Ser Leu Ala Arg Arg Arg
20          25          30

Leu Phe Pro Asn Val Asp Asn Lys Lys Gln Lys Met Ala Gln Pro Gly
35          40          45

Ala Gly Pro Arg Lys Asn Leu Glu Leu Glu Val Met Glu Pro Ala Leu
50          55          60

Lys Asn Asp Gly Pro Ser Leu Asp Thr Ile Leu Val Asn Tyr Leu Asp
65          70          75          80

Thr Leu Thr Gln Arg Val Asn Tyr His Leu Gly Tyr Pro Val Asn Ile
85          90          95

Cys Tyr Asp His Tyr Ala Thr Leu Ala Pro Leu Leu Gln Phe His Leu
100         105         110

```


Asn Asn Cys Gly Asp Pro Phe Leu Gln Asn Thr Val Asp Phe His Ser
 115 120 125
 Lys Asp Phe Glu Val Ala Val Leu Asn Trp Phe Ala Lys Leu Trp Glu
 130 135 140
 Ile Glu Lys Asp Gln Tyr Trp Gly Tyr Val Thr Asn Gly Gly Thr Glu
 145 150 155 160
 Gly Asn Leu His Gly Ile Leu Leu Gly Arg Glu Leu Leu Pro Glu Gly
 165 170 175
 Ile Leu Tyr Ala Ser Lys Asp Ser His Tyr Ser Val Phe Lys Ala Ala
 180 185 190
 Arg Met Tyr Arg Met Asp Ser Glu Thr Ile Asn Thr Ser Val Asn Gly
 195 200 205
 Glu Met Asp Tyr Ser Asp Leu Arg Ala Lys Leu Leu Gln Asn Lys Asp
 210 215 220
 Lys Pro Ala Ile Ile Asn Val Thr Ile Gly Thr Thr Phe Lys Gly Ala
 225 230 235 240
 Ile Asp Asp Leu Asp Val Ile Leu Glu Ile Leu Lys Glu Cys Gly Tyr
 245 250 255
 Ser Gln Asp Arg Phe Tyr Ile His Cys Asp Ala Ala Leu Cys Gly Leu
 260 265 270
 Met Thr Pro Phe Ile Asn Asn Met Ile Ser Phe Lys Lys Pro Ile Gly
 275 280 285
 Ser Val Thr Ile Ser Gly His Lys Phe Leu Gly Cys Pro Met Pro Cys
 290 295 300
 Gly Val Gln Ile Thr Arg Lys Ser Tyr Ile Asn Asn Leu Ser Thr Asn
 305 310 315 320
 Val Glu Tyr Ile Ala Ser Val Asp Ala Thr Ile Ser Gly Ser Arg Asn
 325 330 335
 Gly Leu Thr Pro Ile Phe Leu Trp Tyr Ser Leu Ser Ala Lys Gly Gln
 340 345 350
 Val Gly Leu Gln Lys Asp Val Lys Arg Cys Leu Asp Asn Ala Lys Tyr
 355 360 365
 Leu Lys Asp Arg Leu Gln Gln Ala Gly Ile Ser Val Met Leu Asn Glu
 370 375 380
 Leu Ser Ile Ile Val Val Leu Glu Arg Pro Arg Asp His Glu Phe Val
 385 390 395 400

```
<210> 8
<211> 1416
<212> DNA
<213> Lycopersicon esculentum
```

<400>	8						
atgggtagtc	tctcacttga	aatggatttt	gagccatcac	ctatgacacc	cagaagttta		60
gcagcgatga	cacctagaag	tttagcgcgg	cgaagattgt	ttcccaatgt	ggacaacaaa		120
aaacaaaagg	tgcaacaatc	aggtgcaggg	ccaaggaaga	acttacaact	tgaagtcatg		180
gaacctgcat	tgaacaatgc	tggtccctct	ttggacacta	tattgggtcaa	ttatttagac		240
acatttacac	aacgagtcaa	ttatcattta	ggttatccag	tcaacatttg	ttatgatcac		300
tatgcaactt	tagcaccact	tttacagttt	cacctaaaca	attgtgggtga	tcctttccta		360
caaaacactg	tcgatttcca	ttctaaagac	tttgaagtgg	ctgttttgaa	ttggtttgca		420
aaactatggg	aaattgaaaa	ggatcaatac	tggggatatg	ttaccaatgg	tggcaccgaa		480
ggcaatctcc	atgggtatttt	gttagggaga	gagctacttc	ctgatggaat	attatatgcg		540
tcaaaagact	ctcactattc	ggctcttcaa	gctgcaagaa	tgtatagaat	ggattcagaa		600
acaatcaaca	catcagtaaa	cggagagatg	gattattcag	atttaagagc	aaagttactt		660
caaaataagg	ataaaccagc	tattataaat	gtcacaattg	gaactacgtt	caaaggagca		720
atcgatgacc	tggatgttat	tcttgaaaca	ctcaaagaat	gtggctattc	gcaagatagg		780
ttttacatcc	actgtgatgc	tgcactatgt	ggctcttatga	ccccttttat	aaacaatatg		840
attagtttca	agaagccaat	tggaagtgtc	acaatttctg	gacacaagtt	tttgggatgt		900
ccaatgcctt	gtgggtgtcca	aattacaaga	aagagttaca	tcaataatct	ctcaacaaat		960

```

gtggaataca ttgcttctgt cgatgccact atttctggca gccgtaacgg ttttaactcca 1020
atcttcttgt ggtatagctt gagcgcaaaa ggtcaagttg gacttcaaaa ggatgttaaa 1080
agatgtctcg acaatgccaa atatttgaaa gatcgtcttc aaaaagcagg aataagtgtc 1140
atgttaaatag agcttagcat catagtgtga cttgaaaggc ctcgtgacca tgaatttgtc 1200
cgtcgttggc aactctcatg cgtcaaggat atggcacatg ttattgtaat gccaggcatc 1260
acacgagaaa tgcttgacaa tttcacgagt gaattagtgc aacaaagaaa agtatggtat 1320
caaaatggac agaccaatcc tccttggtgt ggagaggata ttggtgctca aaattgtgca 1380
tgctcttatac ataagattga ctacatctgt ccttag 1416

```

```

<210> 9
<211> 471
<212> PRT
<213> Lycopersicon esculentum
<400> 9

```

```

Met Gly Ser Leu Ser Leu Glu Met Asp Phe Glu Pro Ser Pro Met Thr
1          5          10          15

Pro Arg Ser Leu Ala Ala Met Thr Pro Arg Ser Leu Ala Arg Arg Arg
20          25          30

Leu Phe Pro Asn Val Asp Asn Lys Lys Gln Lys Val Gln Gln Ser Gly
35          40          45

Ala Gly Pro Arg Lys Asn Leu Gln Leu Glu Val Met Glu Pro Ala Leu
50          55          60

Asn Asn Ala Gly Pro Ser Leu Asp Thr Ile Leu Val Asn Tyr Leu Asp
65          70          75          80

Thr Leu Thr Gln Arg Val Asn Tyr His Leu Gly Tyr Pro Val Asn Ile
85          90          95

Cys Tyr Asp His Tyr Ala Thr Leu Ala Pro Leu Leu Gln Phe His Leu
100         105         110

Asn Asn Cys Gly Asp Pro Phe Leu Gln Asn Thr Val Asp Phe His Ser
115         120         125

Lys Asp Phe Glu Val Ala Val Leu Asn Trp Phe Ala Lys Leu Trp Glu
130         135         140

Ile Glu Lys Asp Gln Tyr Trp Gly Tyr Val Thr Asn Gly Gly Thr Glu
145         150         155         160

Gly Asn Leu His Gly Ile Leu Leu Gly Arg Glu Leu Leu Pro Asp Gly

```

S:\Sequences\UF\UF-386CP\UF-386CP-seqlist.doc/DNB/sl

Lys Ile Asp Tyr Ile Cys Pro
465 470

DESCRIPTION

MATERIALS AND METHODS FOR SYNTHESIS OF A FLAVOR AND AROMA VOLATILE IN PLANTS

5

This invention was made with government support under National Science Foundation grant number 7223426-12. The government has certain rights in the invention.

Background of the Invention

10 Fruits are major components of the human diet contributing a large portion of vitamins, minerals, antioxidants, and fiber. While flavor and nutrition composition have clear and profound potential for positive human benefit, they have proven to be difficult traits to modify via either traditional breeding or transgenic approaches due to their generally complex biosynthetic and regulatory pathways. In fact, the biochemical descriptors that comprise flavor
15 are poorly defined. What is typically perceived as flavor in many fruits is the product of a complex interaction among sugars, acids and multiple volatile secondary metabolites (Buttery *et al.*, 1988; Baldwin *et al.*, 2000). Synthesis and accumulation of these compounds is the result of coordinated activity of many genes that may also impact additional aspects of plant growth and development. Effective manipulation of these traits for human benefit will therefore require
20 greater knowledge of the pathways involved and the regulatory systems which control them. Prior to the advent of genomics, researchers could focus on the activity of only one to several genes important in a process of interest and could view their respective effects in relative isolation. From a practical perspective, flavor and nutrition are intimately related and equally important as flavor directly impacts the choice of foods for consumption which, in turn, has
25 positive nutritional consequences on the human diet.

Fruit-bearing crop plants are taxonomically diverse (*e.g.*, pepper, tomato, melons, apples, bananas, grapes). However, they do share common features; most, though not all, fruits are enlarged ovaries. While our knowledge of how domesticated plants came to bear fruit or the mechanisms by which they ripen is still rudimentary, more is known about these processes in
30 tomato (*Lycopersicon esculentum*) than in any other species (see Giovannoni (2001) for review).

Furthermore, a diverse set of Near Isogenic Lines (NILs), single gene ripening mutants, and transgenic lines represent portals through which genetic regulation of fruit development and ripening can be studied (Gray *et al.*, 1994; Giovannoni *et al.*, 1999). The diversity of genetically well characterized tomato germplasm described below (Table 1) is unparalleled in other fruiting species. Nevertheless, it is important to realize that while fruit ripening is a complex sum of coordinately regulated biochemical events that vary from species to species, key regulatory components are likely to be maintained (Hobson and Grierson, 1993). For example, one group has recently identified two genes that are essential for fruit ripening, RIN and NOR (Giovannoni, 2001; Vrebalov *et al.*, 2002). Fruit-specific, ripening-induced homologues of these genes have been identified from strawberry and banana (Vrebalov *et al.*, 2002). Strawberry undergoes a very different ripening program as compared to tomato in that strawberry is non-climacteric (*i.e.* no increase in respiration or ethylene biosynthesis during ripening) and accumulates high levels of anthocyanins rather than carotenoids during fruit maturation. Further, it is anatomically a receptacle, whereas most fruits are ovaries. Banana is interesting in that while its fruit are also expanded carpels, it is a monocot. Apparently similar ripening control shared among monocots and dicots indicates that basic ripening regulation is likely conserved through evolution. In summary, these results suggest that while specific nutritional and flavor components may vary among fruit species they are likely due to regulated metabolic flux through similar pathways with similar genetic control systems. Thus, regulatory and biosynthetic genes identified in tomato will allow for modification of the same or related compounds in a wide range of agriculturally important fruit species.

Tomato has long served as a model system for plant genetics, development, physiology and fruit ripening resulting in the accumulation of substantial information regarding the biology of this economically important plant. Many experimental tools and features of tomato make it ideal for study of fruit ripening; these include extensive germplasm collections, numerous natural, induced, and transgenic mutants, routine transformation technology, a dense and expanding RFLP map, numerous cDNA and genomic libraries, a small genome, relatively short life-cycle and ease of growth and maintenance. In addition, numerous genomic tools that have and continue to be developed include: a) over 140,000 EST sequences (~30,000 non-redundant) from 23 different tomato tissues/treatments (with one-third of the ESTs derived from fruit), b)

EST expression arrays being developed and utilized (see bti.cornell.edu/CGEP/CGEP.html) and c) recent initiation of activities toward development of a tomato physical map anchored to the genetic map to facilitate gene isolation and eventual genome sequencing (Tanksley *et al.*, NSF tomato genome project, 1992). The intense research effort in tomato fruit biology has resulted in many important discoveries that have had a broad impact on the field of plant biology, including control of gene expression by antisense technology, characterization of numerous genes influencing fruit development and ripening, characterization of genes for ethylene synthesis and perception, and the recent connection of ripening regulation and ethylene response to the molecular regulation of floral development (Vrebalov *et al.*, 2002).

Fruit maturation and ripening is the summation of biochemical and physiological changes occurring at the terminal stage of development rendering the organ edible and valuable as an agricultural commodity. These changes frequently include modification of cell wall ultrastructure and texture, conversion of starch to sugars, alterations in pigment and nutrient biosynthesis/accumulation, and heightened levels of flavor and aromatic volatiles (Rhodes, 1980; Hobson and Grierson, 1993). While some ripening effects, such as carotenoid and vitamin C synthesis and accumulation, have direct impact on the nutritive value of mature fruit, others impacting flavor and texture (*e.g.*, volatiles, sugars and acids) can have an indirect impact on human nutrition via their contributions to total consumption levels. In short, "if it tastes better" consumption will increase. This is especially critical as poor food choices exert a disproportional impact on children and members of society on lower rungs of the socio-economic ladder.

Although most fruits display modifications in color, texture, flavor and nutrient composition during maturation, two major classifications of ripening, climacteric and non-climacteric, have been utilized to distinguish fruit on the basis of respiration and ethylene synthesis rates. Climacteric fruits such as tomato, avocado, banana, peaches and apples, are distinguished from non-climacteric fruits such as strawberry, grape and citrus, by their increased respiration and ethylene synthesis rates during ripening (Lelievre *et al.*, 1998). In tomato, ethylene has been shown to be necessary for the coordination and completion of ripening (Yang, 1985; Tucker and Brady, 1987; Klee *et al.*, 1991; Picton *et al.*, 1993; Lanahan *et al.*, 1994). The critical role of ethylene in coordinating climacteric ripening at the molecular level was first

observed via analysis of ethylene inducible ripening-related gene expression in tomato (Lincoln *et al.*, 1987; Maunders *et al.*, 1987). Numerous fruit development-related genes have since been isolated via differential expression patterns and biochemical function (reviewed in Gray *et al.*, 1994). The *in vivo* functions of many fruit development- and ripening-related genes have been tested via antisense repression and/or mutant complementation in tomato. As examples, polygalacturonase was shown to be necessary for ripening-related pectin depolymerization and pathogen susceptibility, yet to have little effect on fruit softening (Smith *et al.*, 1988, Giovannoni *et al.*, 1989, Kramer *et al.*, 1990). Inhibition of phytoene synthase resulted in reduced carotenoid biosynthesis and reduction in fruit and flower pigmentation (Fray and Grierson, 1993). Reduced ethylene evolution resulted in ripening inhibition of ACC synthase (ACS) and ACC oxidase (ACO) antisense lines (Oeller *et al.*, 1991; Hamilton *et al.*, 1990) while introduction of a dominant mutant allele of the *NR* ethylene receptor resulted in plants inhibited in virtually every measurable ethylene response including fruit ripening (Wilkinson *et al.*, 1995; Yen *et al.*, 1995).

Expression analysis of multiple tomato ripening-related genes indicates that a subset exhibit developmentally-controlled ethylene inducibility, *i.e.*, they are ethylene inducible only in ripening fruits. Examples include members of the ACO and ACS gene families (Theologis *et al.*, 1993; Blume and Grierson, 1997; Nakatsuka *et al.*, 1998), the *NR* ethylene receptor (Wilkinson *et al.*, 1995; Payton *et al.*, 1996; Lashbrook *et al.*, 1998) and E8 (Deikman *et al.*, 1992). Additional evidence for non-ethylene mediated ripening control comes from analysis of gene expression in ripening impaired mutants such as *rin* (*ripening-inhibitor*) and *nor* (*non-ripening*) that fail to ripen in response to exogenous ethylene yet display signs of ethylene sensitivity and signaling including induction of some ethylene-regulated genes (Yen *et al.*, 1995). These results suggest that regulatory constraints are placed on climacteric fruit maturation in addition to general ethylene biosynthesis and signaling. Such mechanisms could include fruit-specific regulation of certain subsets of ethylene regulated genes or factors that operate separate from and in addition to ethylene as seems to be the case for both the RIN (Vrebalov *et al.*, 2002) and NOR transcription factors. This is particularly interesting as a greater understanding of the relationship between ethylene, developmental, and environmental signals will likely reveal the impact of various signaling systems on pathways impacting flavor and human nutrition. Indeed numerous environmental factors such as light and temperature can dramatically influence the

degree and rate of fruit ripening with significant impacts on the accumulation of carotenoids and flavor compounds (Hobson and Grierson, 1993; Yen *et al.*, 1997).

Numerous plant metabolites can be listed when the net of “nutritive compounds” is cast. These include various antioxidants, vitamins, minerals, fiber, lipids, and amino acids, to name just a few. In addition, as noted above, one can rationally argue that modification of flavor and additional quality attributes may lead to improved health via increased fruit or vegetable consumption.

Tomato fruits are among the highest source of lycopene, β -carotene, and vitamin C (ascorbate) in the diets of humans in the US, South America, and Europe, with steadily increasing prominence in Asia and the Middle East. In addition to direct nutritive value, carotenoids in particular are metabolized to compounds that impact flavor and aroma of fruit and thus have a significant impact on resulting fresh and processed products. Genes encoding the synthetic steps from phytoene through β -carotene (Bartley *et al.*, 1994; Ronen *et al.*, 1999) are potential regulatory points for modification of carotenoid levels. Indeed, available data indicate that accumulation of lycopene is due to coordinated up-regulation of the genes preceding its synthesis and down-regulation of genes that further metabolize it during ripening (Ronen *et al.*, 1999). Numerous mutant, transgenic, RI and breeding lines that display a wide range of levels of lycopene and β -carotene are available (Table 1). While specific mutants represent some of the catalytic steps (*e.g.*, *r* = phytoene synthase and *cr* and *B* = lycopene cyclase; Hamilton *et al.*, 1990; Ronen *et al.*, 1999) others such as *hp-1* and *hp-2* represent regulators of environmental response. Antisense phytoene synthase tomato lines are greatly reduced in all of the carotenoid-derived volatiles (Baldwin *et al.*, 2000). Furthermore, transgenic and mutant lines altered in ethylene synthesis or perception display variation in carotenoid levels (Table 1).

Table 1. Tomato germplasm altered in carotenoids, flavonoids, vitamin C.

<u>Genotype</u>	<u>Carotenoids</u>	<u>Vit.C</u>	<u>Volatiles</u>	<u>Function</u>
<i>rin</i> ; ripening-inhibitor*	very low	low	NA	MADS-box protein
<i>nor</i> ; non-ripening*	low	low	NA	transcription factor
<i>Nr</i> ; Never-ripe*	low	NA	NA	ethylene receptor
<i>hp-2</i> ; high-pigment-2	high	high	NA	DET1 (light signaling)
<i>cr</i> ; crimson	low B, high L	high	NA	lycopene cyclase
<i>B</i> ; Beta	high B, low L	NA	NA	lycopene cyclase
<i>r</i> ; Phytoene Synthase	low	NA	low	phytoene synthase
<hr/>				
<i>hp-1</i> ; high-pigment-1**	high	high	NA	Not cloned (light signaling)
<i>Nr-2</i> ; Never-ripe-2	low	NA	NA	Not cloned
<i>Gr</i> ; Green-ripe	low	NA	NA	Not cloned
<i>t</i> ; tangerine	low	NA	NA	Not cloned
<i>at</i> ; apricot	low	NA	NA	Not cloned
<i>Cnr</i> ; Clear non-ripening	low	NA	NA	Not cloned
<i>L. esculentum</i> x <i>L. pennellii</i>	low-high	low-high	low-high	
<hr/>				
ACO; ACC oxidase*	low	NA	NA	ethylene Biosynthesis
ACS; ACC synthase*	low	NA	NA	ethylene Biosynthesis
ACD; ACC deaminase*	low	NA	NA	ethylene Biosynthesis
TCTR1; tomato CTR1*	low-high	NA	NA	ethylene signaling MAPKKK

The dashed line separates mutants for which the corresponding gene has been cloned (1st tier) from those which have not (2nd tier). The last tier indicates transgenic lines altered in ethylene synthesis or response and with corresponding changes in carotenoid accumulation. Genotypes indicated with an (*) represent those for which multiple independent transgenic lines are available demonstrating a range of carotenoid accumulation levels. **Three different mutant alleles of *hp-1* each having varying degrees of effect on carotenoid and flavonoid accumulation were provided by M. Koornneef. B = β -carotene. L = lycopene. While quantitative data for vitamin C and volatiles are unknown for many of these lines (NA), their respective phenotypes suggest they are likely to be altered in one or both.

In the case of flavor volatiles, the pathways for synthesis are in many cases not well established. For example, synthesis of apocarotenoids such as β -ionone and β -damascenone is not at all understood. Only recently has an Arabidopsis enzyme, CCD1 (Related to Carotenoid Dioxygenases), that synthesizes apocarotenoids such as β -ionone *in vitro* been identified

(Schwartz *et al.*, 2001). This gene is part of a multigene family, some of which are responsible for synthesis of other apocarotenoids such as ABA (Tan *et al.*, 1997). CCD1 cleaves multiple carotenoid substrates at the 9-10 and 9'-10' bonds, potentially releasing volatiles such as β -ionone, although this has not been established *in vivo*. Similarly, several different volatiles are derived from lipid breakdown (Table 2). The likely first step in their syntheses is the action of a lipoxygenase (LOX) (Riley and Thompson, 1997; Baldwin *et al.*, 2000). Currently there are 14 different EST contigs in the tomato database putatively identified as LOX. Any LOX exhibiting correlation with the lipid-derived volatiles would be a candidate sequence for analyses. It is exactly this sort of correlative biochemical and expression approach that resulted in identification of a key enzyme in strawberry volatile synthesis (Aharoni *et al.*, 2000).

Table 2. The 16 most significant flavor volatiles of tomato

Volatile	Conc. (ppb)	Log odor units	Precursor	Odor Characteristics
<i>cis</i> -3-Hexenal	12,000	3.7	lipid	tomato/green
β -ionone	4	2.8	carotenoid	fruity/floral
Hexanal	3,100	2.8	lipid	green/grassy
β -Damascenone	1	2.7	carotenoid	fruity
1-Penten-3-one	520	2.7	lipid	fruity floral/green
2+3-Methylbutanal	27	2.1	ILE/LEU	musty
<i>trans</i> -2-Hexenal	270	1.2	lipid	green
2-Isobutylthiazole	36	1.0	LEU	tomato vine
1-nitro-2-Phenylethane	17	0.9	PHE	musty, earthy
<i>trans</i> -2-Heptenal	60	0.7	lipid	green
Phenylacetaldehyde	15	0.6	PHE	floral/alcohol
6-Methyl-5-hepten-2-one	130	0.4	carotenoid	fruity, floral
<i>cis</i> -3-Hexenol	150	0.3	lipid	green
2-Phenylethanol	1,900	0.3	PHE	nutty
3-Methylbutanol	380	0.2	LEU	earthy, musty
Methyl salicylate	48	0.08	PHE	wintergreen

Volatiles are ranked by importance based on Odor Units (concentration X humans' ability to detect). Concentrations are average values from typical commercial tomatoes. Odor characteristics were determined by a trained expert panel.

Brief Summary of the Invention

The subject invention concerns polynucleotides encoding a plant 2-phenylethanol dehydrogenase enzyme. In one embodiment, the polynucleotide encodes a tomato 2-phenylethanol dehydrogenase. The subject invention also concerns 2-phenylethanol dehydrogenase polypeptides encoded by polynucleotides of the present invention.

The subject invention also concerns polynucleotides encoding a plant phenylalanine decarboxylase enzyme. In one embodiment, the polynucleotide encodes a tomato phenylalanine decarboxylase. The subject invention also concerns phenylalanine decarboxylase polypeptides encoded by polynucleotides of the present invention.

The subject invention also concerns methods for providing a plant with an increased flavor and aroma volatile. Plants can be transformed with one or more polynucleotides of the present invention. The subject invention also concerns these transformed plant cells, plant tissue, and plants and transgenic progeny thereof.

Brief Description of the Figures

Figures 1A and 1B show the results of GC profiles of volatiles collected from M82 or introgression line 8-2-1 ripe fruit. Introgression line 8-2-1 fruit have higher levels of phenylacetaldehyde and 2-phenylethanol than control M82 fruit.

Figure 2 is a graph of phenylacetaldehyde and 2-phenylethanol levels in control M82 and *L. pennellii* introgression line 8-2-1 fruit. Data are presented as % of control M82 fruit.

Figures 3A and 3B show a full-length cDNA (SEQ ID NO. 1) and amino acid sequence (SEQ ID NO. 2) of a 2-phenylethanol dehydrogenase of the present invention.

Figure 4 shows a pathway for production of the volatiles compounds phenylacetaldehyde and 2-phenylethanol in plants. Phenylalanine is decarboxylated by phenylalanine decarboxylase to form phenethylamine. Phenethylamine is then converted to phenylacetaldehyde by an amine oxidase, followed by conversion to 2-phenylethanol by 2-phenylethanol dehydrogenase.

Figures 5A, 5B, and 5C show the result of alcohol dehydrogenase activities of 2-phenylethanol dehydrogenase on phenylacetaldehyde and related substrates. Activity is determined by the disappearance of substrate and a reduction in OD (340nm). Highest activity levels are observed with phenylacetaldehyde as a substrate.

Figures 6A and 6B show the result of GC profiles of volatiles emitted from wild-type Mitchell Diploid (MD) petunia flowers and transgenic petunia flowers expressing the tomato 2-phenylethanol dehydrogenase gene. In transgenic flowers, higher levels of 2-phenylethanol and lower levels of phenylacetaldehyde are observed as compared to wild-type flowers.

5 **Figure 7** is a graph of the levels of 2-phenylethanol and phenylacetaldehyde emitted from petunia flowers of wild-type Mitchell Diploid (MD) and transgenic lines expressing the tomato 2-phenylethanol dehydrogenase gene.

10 **Figures 8A and 8B** show a full-length coding sequence (SEQ ID NO. 4) and amino acid sequence (SEQ ID NO. 5) of tomato phenylalanine decarboxylase cLEC75E21 of the present invention.

Figures 9A and 9B show a full-length coding sequence (SEQ ID NO. 6) and amino acid sequence (SEQ ID NO. 7) of tomato phenylalanine decarboxylase cLEC73K23 of the present invention.

15 **Figure 10** is a Southern blot showing that cLEC73K23 is present on the *L. pennellii* introgression 8-2-1 on tomato chromosome 8. Restriction patterns of *L. pennellii* and IL8-2-1 genomic DNA hybridized to the cLEC73K23 cDNA are identical, whereas *L. esculentum* M82 restriction patterns are different.

20 **Figures 11A and 11B** show a full-length coding sequence (SEQ ID NO. 8) and amino acid sequence (SEQ ID NO. 9) of an *L. pennellii* phenylalanine decarboxylase similar to cLEC73K23 of the present invention.

Figures 12A and 12B show an amino acid sequence alignment of *L. esculentum* cLEC73K23 (SEQ ID NO. 7), *L. esculentum* cLEC75E21 (SEQ ID NO. 5) and the *L. pennellii* cLEC73K23 homolog (SEQ ID NO. 9).

25 **Figures 13A, 13B, and 13C** show gas chromatography profiles of volatile compounds extracted from *E. coli* cultures expressing the tomato aromatic amino acid decarboxylase genes cLEC73K23 and cLEC75E21. *E. coli* expressing these tomato genes produce phenethylamine in media containing phenylalanine, while control *E. coli* cultures do not.

Brief Description of the Sequences

SEQ ID NO. 1 shows a nucleotide sequence encoding a 2-phenylethanol dehydrogenase according to the present invention.

SEQ ID NO. 2 shows an amino acid sequence of a 2-phenylethanol dehydrogenase encoded by SEQ ID NO. 1 of the present invention.

SEQ ID NO. 3 shows an oligonucleotide PCR primer that can be used according to the present invention.

SEQ ID NO. 4 shows a nucleotide sequence encoding a phenylalanine decarboxylase according to the present invention.

SEQ ID NO. 5 shows an amino acid sequence of a phenylalanine decarboxylase encoded by SEQ ID NO. 4 of the present invention.

SEQ ID NO. 6 shows a nucleotide sequence encoding a phenylalanine decarboxylase according to the present invention.

SEQ ID NO. 7 shows an amino acid sequence of a phenylalanine decarboxylase encoded by SEQ ID NO. 6 of the present invention.

SEQ ID NO. 8 shows a nucleotide sequence encoding a phenylalanine decarboxylase according to the present invention.

SEQ ID NO. 9 shows an amino acid sequence of a phenylalanine decarboxylase encoded by SEQ ID NO. 8 of the present invention.

Detailed Disclosure of the Invention

The subject invention concerns polynucleotides encoding a plant 2-phenylethanol dehydrogenase enzyme. In one embodiment, the polynucleotide encodes a 2-phenylethanol dehydrogenase of tomato. In an exemplified embodiment, the polynucleotide encodes a tomato 2-phenylethanol dehydrogenase polypeptide having an amino acid sequence shown in SEQ ID NO. 2, or an enzymatically functional fragment or variant thereof. In a specific embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 2 comprises the nucleotide sequence shown in SEQ ID NO. 1, or a sequence encoding an enzymatically functional fragment or variant of SEQ ID NO. 2. In one embodiment, the polynucleotide

encoding the amino acid sequence shown in SEQ ID NO. 2 comprises nucleotides 172 to 1155 of the nucleotide sequence shown in SEQ ID NO. 1.

The subject invention also concerns polynucleotides encoding a plant phenylalanine decarboxylase enzyme. In one embodiment, the polynucleotide encodes a phenylalanine decarboxylase of tomato. In an exemplified embodiment, the polynucleotide encodes a tomato phenylalanine decarboxylase polypeptide having an amino acid sequence shown in SEQ ID NO. 5, 7, or 9, or an enzymatically functional fragment or variant thereof. In a specific embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 5, 7, or 9 comprises the nucleotide sequence shown in SEQ ID NO. 4, 6, or 8, or a sequence encoding an enzymatically functional fragment or variant of SEQ ID NO. 5, 7, or 9. In one embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 5 comprises nucleotides 1 to 1395 of the nucleotide sequence shown in SEQ ID NO. 4. In another embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 7 or 9 comprises nucleotides 1 to 1413 of the nucleotide sequence shown in SEQ ID NO. 6 or 8.

The subject invention also concerns polynucleotide expression constructs comprising a polynucleotide sequence of the present invention encoding a plant 2-phenylethanol dehydrogenase. In one embodiment, an expression construct of the invention comprises a polynucleotide sequence encoding a tomato 2-phenylethanol dehydrogenase comprising an amino acid sequence shown in SEQ ID NO. 2, or an enzymatically functional fragment or variant thereof. In a specific embodiment, the polynucleotide sequence comprises a polynucleotide sequence shown in SEQ ID NO. 1, or a sequence encoding an enzymatically functional fragment or variant of SEQ ID NO. 2. In one embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 2 comprises nucleotides 172 to 1155 of the nucleotide sequence shown in SEQ ID NO. 1.

The subject invention also concerns polynucleotide expression constructs comprising a polynucleotide sequence of the present invention encoding a plant phenylalanine decarboxylase. In one embodiment, an expression construct of the invention comprises a polynucleotide sequence encoding a tomato phenylalanine decarboxylase comprising an amino acid sequence shown in SEQ ID NO. 5, 7, or 9, or an enzymatically functional fragment or variant thereof. In a specific embodiment, the polynucleotide sequence comprises a polynucleotide sequence shown

in SEQ ID NO. 4, 6, or 8, or a sequence encoding an enzymatically functional fragment or variant of SEQ ID NO. 5, 7, or 9. In one embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 5 comprises nucleotides 1 to 1395 of the nucleotide sequence shown in SEQ ID NO. 4. In another embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 7 or 9 comprises nucleotides 1 to 1413 of the nucleotide sequence shown in SEQ ID NO. 6 or 8.

Expression constructs of the invention generally include regulatory elements that are functional in the intended host cell in which the expression construct is to be expressed. Thus, a person of ordinary skill in the art can select regulatory elements for use in bacterial host cells, yeast host cells, plant host cells, insect host cells, mammalian host cells, and human host cells. Regulatory elements include promoters, transcription termination sequences, translation termination sequences, enhancers, and polyadenylation elements. As used herein, the term "expression construct" refers to a combination of nucleic acid sequences that provides for transcription of an operably linked nucleic acid sequence. As used herein, the term "operably linked" refers to a juxtaposition of the components described wherein the components are in a relationship that permits them to function in their intended manner. In general, operably linked components are in contiguous relation.

An expression construct of the invention can comprise a promoter sequence operably linked to a polynucleotide sequence encoding a 2-phenylethanol dehydrogenase or phenylalanine decarboxylase of the invention. Promoters can be incorporated into a polynucleotide using standard techniques known in the art. Multiple copies of promoters or multiple promoters can be used in an expression construct of the invention. In a preferred embodiment, a promoter can be positioned about the same distance from the transcription start site in the expression construct as it is from the transcription start site in its natural genetic environment. Some variation in this distance is permitted without substantial decrease in promoter activity. A transcription start site is typically included in the expression construct.

If the expression construct is to be provided in or introduced into a plant cell, then plant viral promoters, such as, for example, a cauliflower mosaic virus (CaMV) 35S (including the enhanced CaMV 35S promoter (see, for example U.S. Patent No. 5,106,739)) or a CaMV 19S promoter can be used. Other promoters that can be used for expression constructs in plants

include, for example, prolifera promoter, Ap3 promoter, heat shock promoters, T-DNA 1'- or 2'- promoter of *A. tumefaciens*, polygalacturonase promoter, chalcone synthase A (CHS-A) promoter from petunia, tobacco PR-1a promoter, ubiquitin promoter, actin promoter, *alcA* gene promoter, *pin2* promoter (Xu *et al.*, 1993), maize *WipI* promoter, maize *trpA* gene promoter (U.S. Patent No. 5,625,136), maize CDPK gene promoter, and RUBISCO SSU promoter (U.S. Patent No. 5,034,322) can also be used. Tissue-specific promoters, for example fruit-specific promoters, such as the E8 promoter of tomato (accession number: AF515784; Good *et al.* (1994)) and the TPRP-F1 promoter (Salts *et al.*, 1991; Santino *et al.*, 1997) can be used. U.S. Patent No. 5,753,475 describes fruit-specific promoter sequences. Flower organ-specific promoters can be used with an expression construct of the present invention for expressing a polynucleotide of the invention in the flower organ of a plant. Examples of flower organ-specific promoters include any of the promoter sequences described in U.S. Patent Nos. 6,462,185; 5,639,948; and 5,589,610. Seed-specific promoters such as the promoter from a β -phaseolin gene (of kidney bean) or a glycinin gene (of soybean), and others, can also be used. Constitutive promoters (such as the CaMV, ubiquitin, actin, or NOS promoter), developmentally-regulated promoters, and inducible promoters (such as those promoters that can be induced by heat, light, hormones, or chemicals) are also contemplated for use with polynucleotide expression constructs of the invention.

For expression in animal cells, an expression construct of the invention can comprise suitable promoters that can drive transcription of the polynucleotide sequence. If the cells are mammalian cells, then promoters such as, for example, actin promoter, metallothionein promoter, NF-kappaB promoter, EGR promoter, SRE promoter, IL-2 promoter, NFAT promoter, osteocalcin promoter, SV40 early promoter and SV40 late promoter, Lck promoter, BMP5 promoter, TRP-1 promoter, murine mammary tumor virus long terminal repeat promoter, STAT promoter, or an immunoglobulin promoter can be used in the expression construct. The baculovirus polyhedrin promoter can be used with an expression construct of the invention for expression in insect cells.

For expression in prokaryotic systems, an expression construct of the invention can comprise promoters such as, for example, alkaline phosphatase promoter, tryptophan (*trp*)

promoter, lambda P_L promoter, β -lactamase promoter, lactose promoter, phoA promoter, T3 promoter, T7 promoter, or tac promoter (de Boer *et al.*, 1983).

Promoters suitable for use with an expression construct of the invention in yeast cells include, but are not limited to, 3-phosphoglycerate kinase promoter, glyceraldehyde-3-phosphate dehydrogenase promoter, metallothionein promoter, alcohol dehydrogenase-2 promoter, and hexokinase promoter.

Expression constructs of the invention may optionally contain a transcription termination sequence, a translation termination sequence, a sequence encoding a signal peptide, and/or enhancer elements. Transcription termination regions can typically be obtained from the 3' untranslated region of a eukaryotic or viral gene sequence. Transcription termination sequences can be positioned downstream of a coding sequence to provide for efficient termination. A signal peptide sequence is a short amino acid sequence typically present at the amino terminus of a protein that is responsible for the relocation of an operably linked mature polypeptide to a wide range of post-translational cellular destinations, ranging from a specific organelle compartment to sites of protein action and the extracellular environment. Targeting gene products to an intended cellular and/or extracellular destination through the use of an operably linked signal peptide sequence is contemplated for use with the polypeptides of the invention. Classical enhancers are cis-acting elements that increase gene transcription and can also be included in the expression construct. Classical enhancer elements are known in the art, and include, but are not limited to, the CaMV 35S enhancer element, cytomegalovirus (CMV) early promoter enhancer element, and the SV40 enhancer element. Intron-mediated enhancer elements that enhance gene expression are also known in the art. These elements must be present within the transcribed region and are orientation dependent. Examples include the maize shrunken-1 enhancer element (Clancy and Hannah, 2002).

DNA sequences which direct polyadenylation of mRNA transcribed from the expression construct can also be included in the expression construct, and include, but are not limited to, an octopine synthase or nopaline synthase signal. The expression constructs of the invention can also include a polynucleotide sequence that directs transposition of other genes, *i.e.*, a transposon.

Expression constructs can also include one or more dominant selectable marker genes, including, for example, genes encoding antibiotic resistance and/or herbicide-resistance for selecting transformed cells. Antibiotic-resistance genes can provide for resistance to one or more of the following antibiotics: hygromycin, kanamycin, bleomycin, G418, streptomycin, paromomycin, neomycin, and spectinomycin. Kanamycin resistance can be provided by neomycin phosphotransferase (NPT II). Herbicide-resistance genes can provide for resistance to phosphinothricin acetyltransferase or glyphosate. Other markers used for cell transformation screening include genes encoding β -glucuronidase (GUS), β -galactosidase, luciferase, nopaline synthase, chloramphenicol acetyltransferase (CAT), green fluorescence protein (GFP), or enhanced GFP (Yang *et al.*, 1996).

The subject invention also concerns polynucleotide vectors comprising a polynucleotide sequence of the invention that encodes a 2-phenylethanol dehydrogenase or phenylalanine decarboxylase. Unique restriction enzyme sites can be included at the 5' and 3' ends of an expression construct or polynucleotide of the invention to allow for insertion into a polynucleotide vector. As used herein, the term "vector" refers to any genetic element, including for example, plasmids, cosmids, chromosomes, phage, virus, and the like, which is capable of replication when associated with proper control elements and which can transfer polynucleotide sequences between cells. Vectors contain a nucleotide sequence that permits the vector to replicate in a selected host cell. A number of vectors are available for expression and/or cloning, and include, but are not limited to, pBR322, pUC series, M13 series, and pBLUESCRIPT vectors (Stratagene, La Jolla, CA).

Polynucleotides of the present invention can be composed of either RNA or DNA. Preferably, the polynucleotides are composed of DNA. The subject invention also encompasses those polynucleotides that are complementary in sequence to the polynucleotides disclosed herein.

Because of the degeneracy of the genetic code, a variety of different polynucleotide sequences can encode 2-phenylethanol dehydrogenase enzymes of the present invention. A table showing all possible triplet codons (and where U also stands for T) and the amino acid encoded by each codon is described in Lewin (1985). In addition, it is well within the skill of a person trained in the art to create alternative polynucleotide sequences encoding the same, or essentially

the same, 2-phenylethanol dehydrogenase or phenylalanine decarboxylase enzymes of the subject invention. These variant or alternative polynucleotide sequences are within the scope of the subject invention. As used herein, references to "essentially the same" sequence refers to sequences which encode amino acid substitutions, deletions, additions, or insertions which do not materially alter the functional activity of the polypeptide encoded by the polynucleotides of the present invention. Allelic variants of the nucleotide sequences encoding a 2-phenylethanol dehydrogenase or phenylalanine decarboxylase of the invention are also encompassed within the scope of the invention.

The subject invention also concerns an isolated plant 2-phenylethanol dehydrogenase. In one embodiment, the 2-phenylethanol dehydrogenase is a 2-phenylethanol dehydrogenase of tomato. In a specific embodiment, the 2-phenylethanol dehydrogenase has an amino acid sequence as shown in SEQ ID NO. 2, or an enzymatically functional fragment or variant thereof. A 2-phenylethanol dehydrogenase enzyme of the invention can be purified using standard techniques known in the art. In one embodiment, a polynucleotide of the invention encoding a 2-phenylethanol dehydrogenase is incorporated into a microorganism such as *E. coli* and the 2-phenylethanol dehydrogenase expressed and then isolated therefrom.

The subject invention also concerns an isolated plant phenylalanine decarboxylase. In one embodiment, the phenylalanine decarboxylase is a phenylalanine decarboxylase of tomato. In a specific embodiment, the phenylalanine decarboxylase has an amino acid sequence as shown in SEQ ID NO. 5, 7, or 9, or an enzymatically functional fragment or variant thereof. A phenylalanine decarboxylase enzyme of the invention can be purified using standard techniques known in the art. In one embodiment, a polynucleotide of the invention encoding a phenylalanine decarboxylase is incorporated into a microorganism such as *E. coli* and the phenylalanine decarboxylase expressed and then isolated therefrom.

Polypeptide fragments according to the subject invention typically comprise a contiguous span of about or at least 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134,

135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 5 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 10 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, or 327 amino acids of SEQ ID NO. 2; or a contiguous span of about or at least 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 15 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 20 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 25 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 30 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406,

407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, or 464 amino acids of SEQ ID NO. 5; or a contiguous span of about or at least 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, or 470 amino acids of SEQ ID NO. 7 or 9.

Polypeptide fragments of the subject invention can be any integer in length from at least about 25 consecutive amino acids to 1 amino acid less than the sequence shown in SEQ ID NO. 2, 5, 7, or 9. Thus, for SEQ ID NO. 2, a polypeptide fragment can be any integer of consecutive

amino acids from about 25 to 327 amino acids; for SEQ ID NO. 5, a polypeptide fragment can be any integer of consecutive amino acids from about 25 to 464 amino acids; for SEQ ID NO. 7 or 9, a polypeptide fragment can be any integer of consecutive amino acids from about 25 to 470 amino acids. The term "integer" is used herein in its mathematical sense and thus representative integers include: 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, and/or 470.

Each polypeptide fragment of the subject invention can also be described in terms of its N-terminal and C-terminal positions. For example, combinations of N-terminal to C-terminal fragments of about 25 contiguous amino acids to 1 amino acid less than the full length

polypeptide of SEQ ID NO. 2, 5, 7, and 9 are included in the present invention. Thus, using SEQ ID NO. 2 as an example, a 25 consecutive amino acid fragment could correspond to amino acids of SEQ ID NO. 2 selected from the group consisting of 1-25, 2-26, 3-27, 4-28, 5-29, 6-30, 7-31, 8-32, 9-33, 10-34, 11-35, 12-36, 13-37, 14-38, 15-39, 16-40, 17-41, 18-42, 19-43, 20-44, 21-45, 22-46, 23-47, 24-48, 25-49, 26-50, 27-51, 28-52, 29-53, 30-54, 31-55, 32-56, 33-57, 34-58, 35-59, 36-60, 37-61, 38-62, 39-63, 40-64, 41-65, 42-66, 43-67, 44-68, 45-69, 46-70, 47-71, 48-72, 49-73, 50-74, 51-75, 52-76, 53-77, 54-78, 55-79, 56-80, 57-81, 58-82, 59-83, 60-84, 61-85, 62-86, 63-87, 64-88, 65-89, 66-90, 67-91, 68-92, 69-93, 70-94, 71-95, 72-96, 73-97, 74-98, 75-99, 76-100, 77-101, 78-102, 79-103, 80-104, 81-105, 82-106, 83-107, 84-108, 85-109, 86-110, 87-111, 88-112, 89-113, 90-114, 91-115, 92-116, 93-117, 94-118, 95-119, 96-120, 97-121, 98-122, 99-123, 100-124, 101-125, 102-126, 103-127, 104-128, 105-129, 106-130, 107-131, 108-132, 109-133, 110-134, 111-135, 112-136, 113-137, 114-138, 115-139, 116-140, 117-141, 118-142, 119-143, 120-144, 121-145, 122-146, 123-147, 124-148, 125-149, 126-150, 127-151, 128-152, 129-153, 130-154, 131-155, 132-156, 133-157, 134-158, 135-159, 136-160, 137-161, 138-162, 139-163, 140-164, 141-165, 142-166, 143-167, 144-168, 145-169, 146-170, 147-171, 148-172, 149-173, 150-174, 151-175, 152-176, 153-177, 154-178, 155-179, 156-180, 157-181, 158-182, 159-183, 160-184, 161-185, 162-186, 163-187, 164-188, 165-189, 166-190, 167-191, 168-192, 169-193, 170-194, 171-195, 172-196, 173-197, 174-198, 175-199, 176-200, 177-201, 178-202, 179-203, 180-204, 181-205, 182-206, 183-207, 184-208, 185-209, 186-210, 187-211, 188-212, 189-213, 190-214, 191-215, 192-216, 193-217, 194-218, 195-219, 196-220, 197-221, 198-222, 199-223, 200-224, 201-225, 202-226, 203-227, 204-228, 205-229, 206-230, 207-231, 208-232, 209-233, 210-234, 211-235, 212-236, 213-237, 214-238, 215-239, 216-240, 217-241, 218-242, 219-243, 220-244, 221-245, 222-246, 223-247, 224-248, 225-249, 226-250, 227-251, 228-252, 229-253, 230-254, 231-255, 232-256, 233-257, 234-258, 235-259, 236-260, 237-261, 238-262, 239-263, 240-264, 241-265, 242-266, 243-267, 244-268, 245-269, 246-270, 247-271, 248-272, 249-273, 250-274, 251-275, 252-276, 253-277, 254-278, 255-279, 256-280, 257-281, 258-282, 259-283, 260-284, 261-285, 262-286, 263-287, 264-288, 265-289, 266-290, 267-291, 268-292, 269-293, 270-294, 271-295, 272-296, 273-297, 274-298, 275-299, 276-300, 277-301, 278-302, 279-303, 280-304, 281-305, 282-306, 283-307, 284-308, 285-309, 286-310, 287-311, 288-312, 289-313, 290-314, 291-315, 292-316, 293-317, 294-318, 295-319, 296-320, 297-321,

298-322, 299-323, 300-324, 301-325, 302-326, 303-327, and 304-328. Similarly, the amino acids corresponding to all other fragments of sizes between 26 consecutive amino acids and 327 consecutive amino acids of SEQ ID NO. 2 are included in the present invention and can also be immediately envisaged based on these examples. Therefore, additional examples, illustrating various fragments of the polypeptides of SEQ ID NO. 2 are not individually listed herein in order to avoid unnecessarily lengthening the specification. Fragment embodiments as described above are also contemplated for the polypeptides of SEQ ID NO. 5, 7, and 9, taking into account that the polypeptides are 465, 471, and 471 amino acids in length, respectively, and are not individually listed herein in order to avoid unnecessarily lengthening the specification.

Polypeptide fragments comprising:

a) 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, or 327 consecutive amino acids of SEQ ID NO. 2;

b) 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117,

118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, and 464 consecutive amino acids of SEQ ID NO. 5; and

c) 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250,

251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, and 470 consecutive amino acids of SEQ ID NO. 7 or 9 may alternatively be described by the formula “n to c” (inclusive), where “n” equals the N-terminal amino acid position and “c” equals the C-terminal amino acid position of the polypeptide. In this embodiment of the invention, “n” is an integer having a lower limit of 1 and an upper limit of the total number of amino acids of the full length polypeptide minus 24 (e.g., 328-24=304 for SEQ ID NO. 2). “c” is an integer between 25 and the number of amino acids of the full length polypeptide sequence (328 for SEQ ID NO. 2) and “n” is an integer smaller than “c” by at least 24. Therefore, for SEQ ID NO. 2, “n” is any integer selected from the list consisting of: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235,

236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, and 304; and “c” is any integer selected from the group consisting of: 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, and 328 provided that “n” is a value less than “c” by at least 24. Every combination of “n” and “c” positions are included as specific embodiments of polypeptide fragments of the invention. All ranges used to describe any polypeptide fragment embodiment of the present invention are inclusive unless specifically set forth otherwise.

Fragments of a plant 2-phenylethanol dehydrogenase or phenylalanine decarboxylase of the invention, as described herein, can be obtained by cleaving the polypeptides of the invention with a proteolytic enzyme (such as trypsin, chymotrypsin, or collagenase) or with a chemical reagent, such as cyanogen bromide (CNBr). Alternatively, polypeptide fragments can be generated in a highly acidic environment, for example at pH 2.5. Polypeptide fragments can also be prepared by chemical synthesis or using host cells transformed with an expression vector comprising a polynucleotide encoding a fragment of a 2-phenylethanol dehydrogenase or

phenylalanine decarboxylase enzyme of the invention, for example, a 2-phenylethanol dehydrogenase that is a fragment of the amino acid sequence shown in SEQ ID NO. 2, or a phenylalanine decarboxylase that is a fragment of the amino acid sequence shown in SEQ ID NO. 5, 7, or 9.

5 Substitution of amino acids other than those specifically exemplified or naturally present in a plant 2-phenylethanol dehydrogenase or phenylalanine decarboxylase enzyme of the invention are also contemplated within the scope of the present invention. For example, non-natural amino acids can be substituted for the amino acids of 2-phenylethanol dehydrogenase or phenylalanine decarboxylase, so long as the 2-phenylethanol dehydrogenase or phenylalanine
10 decarboxylase enzyme having the substituted amino acids retains substantially the same biological activity as the 2-phenylethanol dehydrogenase or phenylalanine decarboxylase in which amino acids have not been substituted. Examples of non-natural amino acids include, but are not limited to, ornithine, citrulline, hydroxyproline, homoserine, phenylglycine, taurine, iodotyrosine, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-aminobutyric acid, 2-amino
15 butyric acid, γ -amino butyric acid, ϵ -amino hexanoic acid, 6-amino hexanoic acid, 2-amino isobutyric acid, 3-amino propionic acid, norleucine, norvaline, sarcosine, homocitrulline, cysteic acid, τ -butylglycine, τ -butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, C-methyl amino acids, N-methyl amino acids, and amino acid analogues in general. Non-natural amino acids also include amino
20 acids having derivatized side groups. Furthermore, any of the amino acids in the protein can be of the D (dextrorotary) form or L (levorotary) form. Allelic variants of a protein sequence of 2-phenylethanol dehydrogenase and phenylalanine decarboxylase enzymes of the present invention are also encompassed within the scope of the invention.

Amino acids can be generally categorized in the following classes: non-polar, uncharged
25 polar, basic, and acidic. Conservative substitutions whereby an 2-phenylethanol dehydrogenase or phenylalanine decarboxylase enzyme of the present invention having an amino acid of one class is replaced with another amino acid of the same class fall within the scope of the subject invention so long as the 2-phenylethanol dehydrogenase or phenylalanine decarboxylase enzyme having the substitution still retains substantially the same biological activity as the 2-
30 phenylethanol dehydrogenase or phenylalanine decarboxylase enzyme that does not have the

substitution. Polynucleotides encoding a 2-phenylethanol dehydrogenase or phenylalanine decarboxylase enzyme having one or more amino acid substitutions in the sequence are contemplated within the scope of the present invention. Table 3 below provides a listing of examples of amino acids belonging to each class.

5

Table 3.	
Class of Amino Acid	Examples of Amino Acids
Nonpolar	Ala, Val, Leu, Ile, Pro, Met, Phe, Trp
Uncharged Polar	Gly, Ser, Thr, Cys, Tyr, Asn, Gln
Acidic	Asp, Glu
Basic	Lys, Arg, His

The subject invention also concerns variants of the polynucleotides of the present invention that encode enzymatically active 2-phenylethanol dehydrogenase or phenylalanine decarboxylase enzymes of the invention. Variant sequences include those sequences wherein one or more nucleotides of the sequence have been substituted, deleted, and/or inserted. The nucleotides that can be substituted for natural nucleotides of DNA have a base moiety that can include, but is not limited to, inosine, 5-fluorouracil, 5-bromouracil, hypoxanthine, 1-methylguanine, 5-methylcytosine, and tritylated bases. The sugar moiety of the nucleotide in a sequence can also be modified and includes, but is not limited to, arabinose, xylulose, and hexose. In addition, the adenine, cytosine, guanine, thymine, and uracil bases of the nucleotides can be modified with acetyl, methyl, and/or thio groups. Sequences containing nucleotide substitutions, deletions, and/or insertions can be prepared and tested using standard techniques known in the art.

Fragments and variants of 2-phenylethanol dehydrogenase and phenylalanine decarboxylase of the present invention can be generated as described herein and tested for the presence of enzymatic function using standard techniques known in the art. For example, for testing fragments and/or variants of a 2-phenylethanol dehydrogenase, the conversion of

phenylacetaldehyde to 2-phenylethanol can be assayed according to the present invention. Thus, an ordinarily skilled artisan can readily prepare and test fragments and variants of a 2-phenylethanol dehydrogenase of the invention and determine whether the fragment or variant retains functional enzymatic activity relative to full-length or a wildtype plant 2-phenylethanol dehydrogenase. Similarly, an assay for the conversion of phenylalanine to phenethylamine can be used to assess enzymatic activity of fragments and/or variants of phenylalanine decarboxylase of the present invention.

Polynucleotides and polypeptides contemplated within the scope of the subject invention can also be defined in terms of more particular identity and/or similarity ranges with those sequences of the invention specifically exemplified herein. The sequence identity will typically be greater than 60%, preferably greater than 75%, more preferably greater than 80%, even more preferably greater than 90%, and can be greater than 95%. The identity and/or similarity of a sequence can be 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99% as compared to a sequence exemplified herein. Unless otherwise specified, as used herein percent sequence identity and/or similarity of two sequences can be determined using the algorithm of Karlin and Altschul (1990), modified as in Karlin and Altschul (1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul *et al.* (1990). BLAST searches can be performed with the NBLAST program, score = 100, wordlength = 12, to obtain sequences with the desired percent sequence identity. To obtain gapped alignments for comparison purposes, Gapped BLAST can be used as described in Altschul *et al.* (1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (NBLAST and XBLAST) can be used. See NCBI/NIH website.

The subject invention also contemplates those polynucleotide molecules having sequences which are sufficiently homologous with the polynucleotide sequences exemplified herein so as to permit hybridization with that sequence under standard stringent conditions and standard methods (Maniatis *et al.*, 1982). As used herein, "stringent" conditions for hybridization refers to conditions wherein hybridization is typically carried out overnight at 20-25 C below the melting temperature (T_m) of the DNA hybrid in 6x SSPE, 5x Denhardt's

solution, 0.1% SDS, 0.1 mg/ml denatured DNA. The melting temperature, T_m , is described by the following formula (Beltz *et al.*, 1983):

$$T_m = 81.5 C + 16.6 \log[Na^+] + 0.41(\%G+C) - 0.61(\% \text{ formamide}) - 600/\text{length of duplex in base pairs.}$$

5 Washes are typically carried out as follows:

(1) Twice at room temperature for 15 minutes in 1x SSPE, 0.1% SDS (low stringency wash).

(2) Once at $T_m - 20^\circ C$ for 15 minutes in 0.2x SSPE, 0.1% SDS (moderate stringency wash).

10 As used herein, the terms "nucleic acid" and "polynucleotide" refer to a deoxyribonucleotide, ribonucleotide, or a mixed deoxyribonucleotide and ribonucleotide polymer in either single- or double-stranded form, and unless otherwise limited, would encompass known analogs of natural nucleotides that can function in a similar manner as naturally-occurring nucleotides. The polynucleotide sequences include the DNA strand
15 sequence that is transcribed into RNA and the strand sequence that is complementary to the DNA strand that is transcribed. The polynucleotide sequences also include both full-length sequences as well as shorter sequences derived from the full-length sequences. Allelic variations of the exemplified sequences also fall within the scope of the subject invention. The polynucleotide sequence includes both the sense and antisense strands either as individual
20 strands or in the duplex.

The subject invention also concerns cells transformed with a polynucleotide of the present invention encoding a 2-phenylethanol dehydrogenase of the invention. In one embodiment, the cell is transformed with a polynucleotide sequence comprising a sequence encoding the amino acid sequence shown in SEQ ID NO. 2, or an enzymatically functional
25 fragment or variant thereof. In a specific embodiment, the cell is transformed with a polynucleotide sequence shown in SEQ ID NO. 1, or a sequence encoding an enzymatically functional fragment or variant of SEQ ID NO. 2. In one embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 2 comprises nucleotides 172 to 1155 of the nucleotide sequence shown in SEQ ID NO. 1.

The subject invention also concerns cells transformed with a polynucleotide of the present invention encoding a phenylalanine decarboxylase of the invention. In one embodiment, the cell is transformed with a polynucleotide sequence comprising a sequence encoding the amino acid sequence shown in SEQ ID NO. 5, 7, or 9, or an enzymatically functional fragment or variant thereof. In a specific embodiment, the cell is transformed with a polynucleotide sequence shown in SEQ ID NO. 4, 6, or 8, or a sequence encoding an enzymatically functional fragment or variant of SEQ ID NO. 5, 7, or 9. In one embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 5 comprises nucleotides 1 to 1395 of the nucleotide sequence shown in SEQ ID NO. 4. In another embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 7 or 9 comprises nucleotides 1 to 1413 of the nucleotide sequence shown in SEQ ID NO. 6 or 8.

Preferably, the polynucleotide sequence is provided in an expression construct of the invention. The transformed cell can be a prokaryotic cell, for example, a bacterial cell such as *E. coli* or *B. subtilis*, or the transformed cell can be a eukaryotic cell, for example, a plant cell, including protoplasts, or an animal cell. Plant cells include, but are not limited to, dicotyledonous, monocotyledonous, and conifer cells. In one embodiment, the plant cell is a cell from tomato. Animal cells include human cells, mammalian cells, avian cells, and insect cells. Mammalian cells include, but are not limited to, COS, 3T3, and CHO cells.

Plants, plant tissues, and plant cells transformed with or bred to contain a polynucleotide of the invention are also contemplated by the present invention. Plants within the scope of the present invention include monocotyledonous plants, such as, for example, rice, wheat, barley, oats, sorghum, maize, sugarcane, pineapple, onion, bananas, coconut, lilies, grasses, and millet. Plants within the scope of the present invention also include dicotyledonous plants, such as, for example, tomato, peas, alfalfa, melon, chickpea, chicory, clover, kale, lentil, soybean, tobacco, potato, sweet potato, radish, cabbage, rape, apple trees, grape, cotton, sunflower, and lettuce; and conifers. Preferably, the plant, plant tissue, or plant cell is tomato. Ornamental and herb plants containing a polynucleotide of the invention are also contemplated within the scope of the invention. Ornamental plants include roses, petunia, carnations, orchids, tulips, and the like. Herb plants include parsley, sage, rosemary, thyme, and the like. Techniques for transforming plant cells with a gene are known in the art and include, for example, *Agrobacterium* infection,

biolistic methods, electroporation, calcium chloride treatment, *etc.* Transformed cells can be selected, redifferentiated, and grown into plants using standard methods known in the art. The seeds and progeny of any transformed or transgenic plant cells or plants of the invention are also included within the scope of the present invention.

5 The subject invention also concerns methods for providing a plant with increased flavor or fragrance of fruit or flower by incorporating one or more polynucleotide of the present invention in the genome of the plant cells and expressing the polypeptide encoded by the polynucleotide. In one embodiment, a plant is grown from a transformed plant cell of the invention. Preferably, the polynucleotide encodes a 2-phenylethanol dehydrogenase or a
10 phenylalanine decarboxylase derived from the same plant species as the plant. In one embodiment, the plant is tomato. In another embodiment, the plant is a rose or other scented ornamental. In those embodiments, where increased flavor of fruit is desired, preferably the polynucleotide(s) of the invention is expressed in the fruit. In those embodiments where increased or enhanced fragrance of fruit or flower is desired, preferably the polynucleotide(s) of
15 the invention is expressed in the fruit and/or flower. In a specific embodiment, a polynucleotide encoding an amino acid sequence shown in SEQ ID NO. 2, or an enzymatically functional fragment or variant thereof, is incorporated into a tomato plant genome. In a specific embodiment, the polynucleotide comprises a nucleotide sequence shown in SEQ ID NO. 1, or a sequence encoding an enzymatically functional fragment or variant of SEQ ID NO. 2. In one
20 embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 2 comprises nucleotides 172 to 1155 of the nucleotide sequence shown in SEQ ID NO. 1. In another specific embodiment, a polynucleotide encoding an amino acid sequence shown in SEQ ID NO. 5, 7, or 9, or an enzymatically functional fragment or variant thereof, is incorporated into a tomato plant genome. In a specific embodiment, the polynucleotide comprises a nucleotide
25 sequence shown in SEQ ID NO. 4, 6, or 8, or a sequence encoding an enzymatically functional fragment or variant of SEQ ID NO. 5, 7, or 9. In one embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 5 comprises nucleotides 1 to 1395 of the nucleotide sequence shown in SEQ ID NO. 4. In another embodiment, the polynucleotide encoding the amino acid sequence shown in SEQ ID NO. 7 or 9 comprises nucleotides 1 to 1413.
30 of the nucleotide sequence shown in SEQ ID NO. 6 or 8. The level of expression of a

polynucleotide of the invention can be manipulated using standard methods known in the art, including the use of promoters that provide for low, intermediate or high levels of expression.

The subject invention also concerns methods for producing 2-phenylethanol. In one embodiment, recombinantly produced 2-phenylethanol dehydrogenase of the invention can be used to enzymatically convert a suitable substrate, such as phenylacetaldehyde, into 2-phenylethanol. In another embodiment, a microorganism, such as yeast or *E. coli*, can be transformed with and express a polynucleotide encoding a plant 2-phenylethanol dehydrogenase enzyme of the invention and, optionally, one or more enzymes, such as phenylalanine decarboxylase and phenylethylamine oxidase, that through their enzymatic reactions result in a suitable substrate (*e.g.*, phenylacetaldehyde) for 2-phenylethanol dehydrogenase to convert to 2-phenylethanol. Transformed microorganisms can be grown and polynucleotides expressed constitutively or induced, and 2-phenylethanol isolated from the microorganisms.

The subject invention also concerns methods for producing phenethylamine. In one embodiment, recombinantly produced phenylalanine decarboxylase of the invention can be used to enzymatically convert a suitable substrate, such as phenylalanine, into phenethylamine. In another embodiment, a microorganism, such as yeast or *E. coli*, can be transformed with and express a polynucleotide encoding a plant phenylalanine decarboxylase of the invention and, optionally, one or more enzymes, that through their enzymatic reactions result in a suitable substrate (*e.g.*, phenylalanine) for phenylalanine decarboxylase to convert to phenethylamine. Transformed microorganisms can be grown and polynucleotides expressed constitutively or induced, and phenethylamine isolated from the microorganisms.

All patents, patent applications, provisional applications, and publications referred to or cited herein are incorporated by reference in their entirety, including all figures and tables, to the extent they are not inconsistent with the explicit teachings of this specification.

Materials and Methods

Plant material.

Tomato (*Lycopersicon esculentum* Mill. cv. M82) and *Lycopersicon pennellii* introgression lines 8-2 and 8-2-1 (Eshed and Zamir, 1994) were grown in the greenhouse or field

under standard conditions. Petunia plants were grown in a greenhouse under standard conditions.

Volatile collection.

Volatiles were collected from approximately 100g of chopped ripe tomato fruit as described by Schmelz *et al.* (2003). Petunia volatiles were collected from five flowers from each plant harvested at dusk. Volatiles were separated on an Agilent DB-5 column and analyzed on an Agilent 6890N gas chromatograph.

Microarrays.

Tomato cDNA microarrays were as described in Moore *et al.* (2002). Total RNAs were isolated as described earlier (Ciardi *et al.*, 2000). Arrays were hybridized with Cy3 or Cy5 labeled cDNAs from M82 and introgression line 8-2-1 fruit. Arrays were performed multiple times and with dyes reversed to ensure accuracy of the expression data.

2-phenylethanol dehydrogenase expression in *E. coli*.

A full-length 2-phenylethanol dehydrogenase cDNA was cloned by 5' RACE from tomato fruit cDNA using primer 5'-TCCTTGCCCCACCAAGAGAAAGCAAGTGCTGCGT-3' (SEQ ID NO. 3). Following sequence analysis the full-length cDNA was obtained by PCR.

The coding region was cloned into vector pDEST15 containing a glutathione S-transferase (GST) tag (Invitrogen) by recombination, and transformed into *E. coli* strain BL21-SI (Invitrogen) for inducible protein expression. Enzyme activity of crude *E. coli* extracts was determined by the method of Larroy *et al.* (2002) using phenylacetaldehyde, cinnamaldehyde or vanillin as a substrate.

Production of transgenic petunia plants.

The full-length 2-phenylethanol dehydrogenase cDNA (SEQ ID NO. 1) was cloned in a vector under the control of the figwort mosaic virus 35S promoter (Richins *et al.*, 1987) and followed by the *Agrobacterium* nopaline synthase (*nos*) 3' terminator. The transgene was

introduced into *Petunia hybrida* cv. Mitchell Diploid by the method of Wilkinson *et al.* (1997) with kanamycin resistance as a selectable marker.

In vivo phenylalanine decarboxylase activity assays.

5 Tomato (M82 or introgression line 8-2-1) fruit pericarp disks were incubated with 1μCi universally-labeled ¹⁴C-phenylalanine for 8hr. Production of ¹⁴C-CO₂ was measured by incubating the pericarp disk in a sealed flask with a 2N KOH filter paper disk suspended above the pericarp disk, followed by scintillation counting. ¹⁴C-phenylalanine and ¹⁴C-phenylethylamine were extracted from the pericarp disk and separated using an AG-1 (OH-) 10 column in series with a BioRex-70 column as described by Rontein *et al.* (2001). Production of ¹⁴C-phenylethylamine was confirmed by thin layer chromatography.

Phenylalanine decarboxylase expression in *E. coli*.

The full-length aromatic amino acid decarboxylase cDNAs were cloned by sequencing 15 putative clones from the TIGR database. Following sequence analysis the full-length coding sequence was obtained by PCR, and cloned into vector pENTR/D-TOPO. The coding region was then cloned into vector pDEST15 containing a GST tag (Invitrogen) by recombination, and transformed into *E. coli* strain BL21-AI (Invitrogen) for inducible protein expression. Production of the recombinant protein was confirmed by protein blotting with anti-GST 20 antibodies. Enzyme activity was determined by growing *E. coli* strains expressing the aromatic amino acid decarboxylases in media containing 19.4mM phenylalanine. Volatile compounds were extracted from the cultures using an equal volume of hexanes. Extracts were concentrated and analyzed by gas chromatography on an Agilent DB-5 column on an Agilent 6890N gas chromatograph. Identification of phenethylamine was confirmed by GC-mass spectrometry as 25 described by Schmelz *et al.* (2001).

Southern blotting.

DNA from *L. esculentum* M82 or 8-2-1 or *L. pennellii* leaves was digested with *EcoRI*, 30 *EcoRV*, *DraI*, *HaeIII* or *ScaI*. Southern blotting was performed as described by Sambrook *et al.* (1989).

Following are examples which illustrate procedures for practicing the invention. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted.

5

Example 1—Volatiles Analysis

Volatiles analysis by GC on a DB5 column indicated high levels of phenylacetaldehyde and 2-phenylethanol in ripe fruit from *L. pennellii* introgression line (IL) 8-2-1. Levels of other tomato volatiles were similar to control M82 ripe fruit (Figure 1). Levels of 2-phenylethanol
10 were approximately 250X higher in 8-2-1 fruit than in M82 fruit. Phenylacetaldehyde levels in IL8-2-1 fruit were approximately 20X higher than in control fruit (Figure 2). IL8-2-1 fruit had a distinct floral (rose-like) aroma consistent with the floral aromas of phenylacetaldehyde and 2-phenylethanol.

Example 2—Microarray Analysis and Isolation of 2-Phenylethanol Dehydrogenase

15 Microarrays containing approximately 3,000 tomato cDNAs indicated that an alcohol dehydrogenase gene (clone cLET2M9) was more highly expressed in IL8-2-1 fruit than in control M82 fruit; whereas, a related tomato alcohol dehydrogenase gene was not upregulated in IL8-2-1 fruit (Table 4).

Table 4. Microarray gene expression data for two alcohol dehydrogenase-like genes	
Microarray clone	Ratio
cLET2M9	+2.2
cLEG71F12	-2.35

20 RNAs extracted from M82 and introgression line 8-2-1 fruit were compared using cDNA microarrays. Positive values indicate higher RNA expression levels in introgression line 8-2-1 fruit; negative values indicate higher expression in M82 fruit.

25 Since clone cLET2M9 was only a partial cDNA, the full-length cDNA sequence for this clone was obtained by 5' RACE. The full-length cDNA sequence of 2-phenylethanol dehydrogenase was then obtained by PCR, and confirmed by sequence analysis (Figure 3A) (SEQ ID NO. 1). The deduced amino acid sequence of the 2-phenylethanol dehydrogenase is

shown in Figure 3B (SEQ ID NO. 2). In plants, a substrate such as phenylacetaldehyde can be converted to 2-phenylethanol by 2-phenylethanol dehydrogenase (Figure 4).

Example 3—Enzyme Activity

5 The 2-phenylethanol dehydrogenase coding region of the full-length cLET2M9 and the related cLEG71F12 were cloned in vector pDEST15 with a GST tag and transformed into *E. coli* BL21-SI cells for inducible expression. The production of recombinant protein in *E. coli* was determined by Western blotting with an anti-GST antibody. Alcohol dehydrogenase activities on phenylacetaldehyde and several related substrates were determined spectrophotometrically by
10 the reduction in levels of NADPH and a decrease in OD₃₄₀ (Figure 5). The highest level of activity was observed with phenylacetaldehyde as a substrate. Lower activity levels were also observed with cinnamaldehyde as a substrate, whereas negligible activity was seen with vanillin as a substrate. Protein from cLEG71F12 expressed in *E. coli* showed very little activity on the three substrates tested.

Example 4—Expression of 2-Phenylethanol Dehydrogenase in Transgenic Petunia

15 Full-length tomato 2-phenylethanol dehydrogenase cDNA was introduced into petunia (cv. Mitchell Diploid) under control of the constitutively expressed figwort mosaic virus promoter. Several transgenic petunia lines had high levels of expression of the tomato gene in
20 flowers (data not shown). Wild-type petunia flowers emit relatively high levels of phenylacetaldehyde and lower levels of 2-phenylethanol. However, the transgenic petunia flowers expressing the polynucleotide encoding tomato 2-phenylethanol dehydrogenase have higher levels of 2-phenylethanol and lower levels of phenylacetaldehyde than wild-type flowers (Figure 6). Levels of other petunia flower volatiles were similar to wild-type in the transgenic
25 flowers. A range of phenylacetaldehyde and 2-phenylethanol levels were seen in the transgenic lines, however the majority of the lines had higher levels of 2-phenylethanol and lower levels of phenylacetaldehyde than wild-type flowers (Figure 7). Overall, these data indicate that the introduction of the 2-phenylethanol dehydrogenase tomato transgene results in the conversion of phenylacetaldehyde to 2-phenylethanol in petunia flowers.

Example 5—*In vivo* Aromatic Amino Acid Decarboxylase Activity

A pathway for the formation of 2-phenylethanol in plants is shown in Figure 4. In the pathway phenylalanine is converted to phenethylamine by phenylalanine decarboxylase, followed by conversion of phenethylamine to phenylacetaldehyde by an amine oxidase. Phenylacetaldehyde is then converted to 2-phenylethanol by a 2-phenylethanol dehydrogenase. To establish the first step in this pathway, tomato fruit pericarp disks were incubated with ^{14}C -phenylalanine. The formation of ^{14}C -phenethylamine and ^{14}C - CO_2 , the products of the decarboxylase reaction were then determined (Table 5). Higher levels of both phenethylamine and CO_2 were formed in the IL8-2-1 pericarp disks than in the control M82 pericarp disks (Table 5), and correlate with the higher levels of phenylacetaldehyde and 2-phenylethanol in IL8-2-1 tomato fruit. These results indicate that phenethylamine can be an intermediate in the pathway to phenylacetaldehyde and 2-phenylethanol.

Table 5. Phenylalanine decarboxylase activity of M82 and IL8-2-1 tomato pericarp disks fed ^{14}C phenylalanine.

Line	nCi CO_2	nCi phenethylamine
M82	0.96 ± 0.05	0.36 ± 0.02
8-2-1	5.16 ± 2.36	2.08 ± 0.39

Pericarp disks were fed 1 μCi universally labeled ^{14}C -phenylalanine for 8 hours, and amounts of ^{14}C - CO_2 and ^{14}C -phenethylamine produced were determined.

Example 6—Identification of Aromatic Amino Acid Decarboxylase Genes

Conversion of phenylalanine to phenethylamine would be catalyzed by a phenylalanine decarboxylase. Therefore, tomato sequence databases were searched for cDNAs with similarity to other aromatic amino acid decarboxylases. Several clones of *L. esculentum* cDNA sequences were identified, although two were more similar to histidine decarboxylases than aromatic amino acid decarboxylases. The full-length coding sequence of each of these genes was obtained by PCR, and confirmed by sequence analysis.

Example 7—Enzyme Activity

The coding regions of the full-length aromatic amino acid decarboxylases were cloned in vector pDEST15 with a GST tag and transformed into *E.coli* BL21-SI cells for inducible expression. The production of recombinant proteins in *E. coli* was confirmed by Western blotting with an anti-GST antibody. *E. coli* cultures expressing the putative aromatic amino acid decarboxylases were grown in the presence of phenylalanine. Volatile compounds were extracted from these cultures and analyzed by gas chromatography. *E. coli* cultures expressing two putative aromatic amino acid decarboxylases (cLEC73K23 and cLEC75E21) produced a compound with the same retention time as phenethylamine, while control cultures did not (Figure 13). *E. coli* cultures expressing three other putative amino acid decarboxylases did not produce phenethylamine. The presence of phenethylamine in the samples was confirmed by GC-MS. The nucleic acid coding sequences and amino acid sequences of the phenylalanine decarboxylases are shown in Figures 8A-8B (SEQ ID NO. 4 and SEQ ID NO. 5) and Figures 9A-9B (SEQ ID NO. 6 and SEQ ID NO. 7). The cLEC73K23 and cLEC75E21 cDNA sequences exhibit 79% identity to one another, whereas the amino acid sequences are 85% similar and 81% identical to each other.

Example 8—Southern Blotting

To determine if either of the genes with phenylalanine decarboxylase activity could be responsible for the altered levels of phenylalanine-derived volatiles in 8-2-1 fruit with the *L. pennellii* introgression on chromosome 8, Southern blots on M82, 8-2-1 and *L. pennellii* genomic DNA were performed using cLEC73K23 as a probe. Polymorphisms between M82 and *L. pennellii* were observed with *EcoRI*, *EcoRV* and *ScaI* (Figure 10). With each of these enzymes the 8-2-1 restriction pattern was identical to *L. pennellii*, indicating that the cLEC73K23 gene in introgression line 8-2-1 came from the *L. pennellii* parent.

Example 9—Cloning of the *L. pennellii* cLEC73K23 Sequence

The nucleic acid coding sequence of the *L. pennellii* cLEC73K23 gene was obtained by PCR with primers from the 5' and 3' ends of the *L. esculentum* cLEC73K23. The correct 5' and 3' ends of the *L. pennellii* gene were then obtained by 5' and 3' RACE. The full-length coding

sequence of this gene and the amino acid sequence are shown in Figures 11A–11B (SEQ ID NO. 8 and SEQ ID NO. 9). The *L. pennellii* and *L. esculentum* cLEC73K23 cDNA sequences are 95% identical, whereas the amino acid sequences are 98% similar and 97% identical. An amino acid alignment of the *L. pennellii* and *L. esculentum* cLEC73K23 sequences and the *L.*
5 *esculentum* cLEC75E21 sequences are shown in Figure 12.

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this
10 application. In addition, any elements or limitations of any invention or embodiment thereof disclosed herein can be combined with any and/or all other elements or limitations (individually or in any combination) or any other invention or embodiment thereof disclosed herein, and all such combinations are contemplated with the scope of the invention without limitation thereto.

References

- U.S. Patent No. 5,106,739
- 5 U.S. Patent No. 5,625,136
- U.S. Patent No. 5,034,322
- U.S. Patent No. 5,589,610
- 10 U.S. Patent No. 5,639,948
- U.S. Patent No. 5,753,475
- 15 U.S. Patent No. 6,462,185
- Aharoni, A. *et al.* (2000) "Identification of the SAAT gene involved in strawberry flavor biogenesis by use of DNA microarrays" *Plant Cell* 12:647-661.
- 20 Altschul, S. F. *et al.* (1990) "Basic Local Alignment Search Tool" *J. Mol. Biol.* 215:402-410.
- Altschul, S. F. *et al.* (1997) "Gapped BLAST and PSI-BLAST: A New Generation of Protein Database Search Programs" *Nucl. Acids Res.* 25:3389-3402.
- 25 Baldwin, E.A. *et al.* (2000) "Flavor trivia and tomato aroma: Biochemistry and possible mechanisms for control of important aroma components" *Hortscience* 35:1013-1022.
- Bartley, G. *et al.* (1994) "Molecular biology of carotenoid biosynthesis in plants" *Annual Review of Plant Physiology & Plant Molecular Biology* Jones, R. L.; Somerville, C. R.: Eds.
- 30 Annual Review of Plant Physiology and Plant Molecular Biology. 45: 287-301.
- Beltz, G. A., Jacobs, K. A., Eickbush, T. H., Cherbas, P. T., Kafatos, F. C. (1983) "Isolation of multigene families and determination of homologies by filter hybridization methods" *Methods of Enzymology*, R. Wu, L. Grossman and K. Moldave [eds.] Academic Press,
- 35 New York 100:266-285.
- Blume, B. and Grierson, D. (1997) "Expression of ACC oxidase promoter-GUS fusions in tomato and *Nicotiana plumbaginifolia* regulated by developmental and environmental stimuli" *Plant J.* 12: 731-746.
- 40 Buttery, R. *et al.* (1988) "Quantitative studies on origins of fresh tomato aroma volatiles" *J. Agric. Food Chem.* 36:1247-1250.

- Ciardi, J.A. *et al.* (2000) "Response to *Xanthomonas campestris* pv. *vesicatoria* in tomato involves regulation of ethylene receptor gene expression" *Plant Physiol.* 123:81-92.
- 5 Clancy, M. and Hannah, L.C. (2002) "Splicing of the maize Sh1 first intron is essential for enhancement of gene expression, and a T-rich motif increases expression without affecting splicing," *Plant Physiol.* 130(2):918-29.
- 10 de Boer, H. A., Comstock, L. J., Vasser, M. (1983) "The tac promoter: a functional hybrid derived from the trp and lac promoters" *Proc. Natl. Acad. Sci. USA* 80(1):21-25.
- Deikman, J. *et al.* (1992) "Organization of ripening and ethylene regulatory regions in a fruit-specific promoter from tomato (*Lycopersicon esculentum*)" *Plant Physiol.* 100:2013-2017.
- 15 Eshed, Y. and Zamir, D. (1994) "A genomic library of *Lycopersicon pennellii* in *L. esculentum*: A tool for fine mapping of genes" *Euphytica* 79:175-179.
- 20 Fray, R. and Grierson, D. (1993) "Identification and genetic analysis of normal and mutant phytoene synthase genes of tomato by sequencing, complementation, and co-suppression" *Plant Mol. Biol.* 22:589-602.
- Giovannoni, J. (2001) "Molecular Regulation of Fruit Ripening" *Ann. Rev. Plant Physiol. Plant Molec. Biol.* 52:725-749.
- 25 Giovannoni, J. *et al.* (2002) "A MADS-box gene necessary for fruit ripening at the tomato ripening-inhibitor (*rin*) locus" *Science* 296(5566):275-276.
- 30 Giovannoni, J. *et al.* (1999) "Analysis of gene expression and mutants influencing ethylene responses and fruit development in tomato" In Biology and Biotechnology of the Plant Hormone Ethylene II, (Kanellis, A. ed.) Kluwer Academic Publishers, pp. 119-127.
- 35 Giovannoni, J. *et al.* (1989) "Expression of a chimeric polygalacturonase gene in transgenic *rin* (ripening inhibitor) tomato fruit results in polyuronide degradation but not fruit softening" *Plant Cell* 1:53-63.
- Good, X. *et al.* (1994) "Reduced ethylene synthesis by transgenic tomatoes expressing S-adenosylmethionine hydrolase." *Plant Molec. Biol.* 26:781-790.
- 40 Gray, J.E. *et al.* (1994) "The use of transgenic and naturally occurring mutants to understand and manipulate tomato fruit ripening" *Plant, Cell and Environment* 17:557-571.
- Hamilton, A. *et al.* (1990) "Antisense gene that inhibits synthesis of the hormone ethylene in transgenic plants" *Nature* 346:284-287.

- Hobson, G. and Grierson, D. (1993) "Tomato" In: Seymour GB, Taylor JE, Tucker GA (eds) *Biochemistry of Fruit Ripening*. Chapman and Hall, London pp 405-442.
- 5 Karlin S. and Altschul, S. F. (1990) "Methods for Assessing the Statistical Significance of Molecular Sequence Features by Using General Scoring Schemes" *Proc. Natl. Acad. Sci. USA* 87:2264-2268.
- 10 Karlin S. and Altschul, S. F. (1993) "Applications and Statistics for Multiple High-Scoring Segments in Molecular Sequences" *Proc. Natl. Acad. Sci. USA* 90:5873-5877.
- Klee, H.J. *et al.* (1991) "Control of ethylene synthesis by expression of a bacterial enzyme in transgenic tomato plants" *Plant Cell* 3:1187-1193.
- 15 Kramer, M. *et al.* (1990) "Field evaluation of tomatoes with reduced polygalacturonase by antisense RNA" In *Horticultural Biotechnology*. Bennett, A. and O'Neill, S. (eds.) Alan R. Liss. pp. 347-355.
- 20 Lanahan, M.B. *et al.* (1994) "The Never Ripe mutation blocks ethylene perception in tomato" *Plant Cell* 6:521-530.
- Larroy, C. *et al.* (2002) "Characterization of the *Saccharomyces cerevisiae* YMR318C (*ADH6*) gene product as a broad specificity NADPH-dependent alcohol dehydrogenase: relevance in aldehyde reduction" *Biochem. J.* 361:163-172.
- 25 Lashbrook, C. *et al.* (1998) "Transgenic analysis of tomato endo-beta-1,4-glucanase gene function. Role of CEL1 in floral abscission" *Plant J.* 13:303-310.
- Lelievre, J. *et al.* (1998) "Ethylene and fruit ripening" *Plant Physiol.* 101:727-739.
- 30 Lewin, B. (1985) *Genes II*, John Wiley & Sons, Inc., p. 96.
- Lincoln, J. *et al.* (1987) "Regulation of gene expression by ethylene during *Lycopersicon esculentum* (tomato) fruit development" *Proc. Nat. Acad. Sci. USA*. 84:2793-2797.
- 35 Maniatis, T., E.F. Fritsch, J. Sambrook (1982) "Nuclease *Bal31*" *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- 40 Maunders, M. *et al.* (1987) "Ethylene stimulates the accumulation of ripening-related mRNAs in tomatoes" *Plant Cell Environ.* 10:177-184.
- Moore, S. *et al.* (2002) "Use of genomics tools to isolate key ripening genes and analyse fruit maturation in tomato" *J. Exp. Bot.* 53:2023-2030.
- 45 Nakatsuka, A. *et al.* (1998) "Differential expression and internal feedback regulation of 1-aminocyclopropane-1-carboxylate synthase, of 1-aminocyclopropane-1-carboxylate

- Oeller, P.W. *et al.* (1991) "Reversible inhibition of tomato fruit senescence by antisense 1-aminocyclopropane-1-carboxylate synthase" *Science* 254:427-439.
- 5 Payton, S. *et al.* (1996) "Ethylene receptor expression is regulated during fruit ripening, flower senescence and abscission" *Plant Molecular Biology* 31:1227-1231.
- Picton, S. *et al.* (1993) "Altered fruit ripening and leaf senescence in tomatoes expressing an antisense ethylene-forming enzyme transgene" *Plant J.* 3:469-481.
- 10 Rhodes, M.J.C. (1980) "The maturation and ripening of fruits" In: KV Thimann (ed) *Senescence in Plants*. CRC Press, Boca Raton, FL, pp. 157-205.
- Richins, R.D. *et al.* (1987) Sequence of the figwort mosaic virus DNA (caulimovirus group)" *Nucl. Acids Res.* 15:8451-8466.
- 15 Riley, J. and Thompson, J. (1997) "Subcellular generation and distribution of lipid-derived volatiles in the ripe tomato" *J. Plant Physiol.* 149-150:546-551.
- Ronen, G. *et al.* (1999) "Regulation of carotenoid biosynthesis during tomato fruit development: Expression of the gene for lycopene epsilon-cyclase is down-regulated during ripening and is elevated in the mutant Delta" *Plant J.* 17: 341-351.
- 20 Rontein, D. *et al.* (2001) "Plants synthesize ethanolamine by direct decarboxylation of serine using a pyridoxal phosphate enzyme" *J. Biol. Chem.* 276:35523-35529.
- 25 Sambrook, J. *et al.* (1989) Molecular Cloning: A Laboratory Manual, 2nd ed. (Cold Spring Harbor, N.Y., Cold Spring Harbor Laboratory).
- Schmelz, E.A. *et al.* (2001) "The influence of intact-plant and excised-leaf bioassay designs on volicitin- and jasmonic acid-induced sesquiterpene volatile release in *Zea mays*" *Planta* 214:171-179.
- 30 Schmelz, E.A. *et al.* (2003) "Qualitative relationships between induced jasmonic acid levels and volatile emission in *Zea mays* during *Spodoptera exigua* herbivory" *Planta* 216:665-673.
- 35 Schwartz, S. *et al.* (2001) "Characterization of a novel carotenoid cleavage dioxygenase from plants" *J. Biol. Chem.* 276:25208-25211.
- Smith, C. *et al.* (1988) "Antisense RNA inhibition of polygalacturonase gene expression in transgenic tomatoes" *Nature* 334:724-726.
- 40 Tan, B. *et al.* (1997) "Genetic Control of Absciscic Acid Biosynthesis in Maize" *Proc. Nat. Acad. Sci. USA* 94:12235-12240.

- Schwartz, S. *et al.* (2001) "Characterization of a novel carotenoid cleavage dioxygenase from plants" *J. Biol. Chem.* 276:25208-25211.
- 5 Smith, C. *et al.* (1988) "Antisense RNA inhibition of polygalacturonase gene expression in transgenic tomatoes" *Nature* 334:724-726.
- Tan, B. *et al.* (1997) "Genetic Control of Absciscic Acid Biosynthesis in Maize" *Proc. Nat. Acad. Sci. USA* 94:12235-12240.
- 10 Tanksley, S. *et al.* (1992) "High density molecular linkage maps of the tomato and potato genomes" *Genetics* 132:1141-1160.
- Theologis, A. (1993) "Use of a tomato mutant constructed with reverse genetics to study fruit ripening, a complex developmental process" *Dev. Genetics* 14:282-259.
- 15 Tucker, G.A. and Brady, C.J. (1987) "Silver ions interrupt tomato fruit ripening" *J. Plant Physiol.* 127:165-169.
- Vrebalov, J. *et al.* (2002) "A MADS-box gene necessary for fruit ripening at the tomato ripening-inhibitor (*rin*) locus" *Science* 296:343-346.
- 20 Wilkinson, J.Q. *et al.* (1997) "A dominant mutant receptor from *Arabidopsis* confers ethylene insensitivity in heterologous plants" *Nature Biotechnol.* 15: 444-447.
- 25 Wilkinson, J. *et al.* (1995) "An ethylene-inducible component of signal transduction encoded by Never-ripe" *Science* 270:1807-1809.
- Xu, D., McElroy, D., Thornburg, R. W., Wu, R. *et al.* (1993) "Systemic induction of a potato *pin2* promoter by wounding, methyl jasmonate, and absciscic acid in transgenic rice plants" *Plant Molecular Biology* 22:573-588.
- 30 Yang, S.F. (1985) "Biosynthesis and action of ethylene" *HortScience* 20:41-45.
- Yang, T. T. *et al.* (1996) "Optimized Codon Usage and Chromophore Mutations Provide Enhanced Sensitivity with the Green Fluorescent Protein" *Nucleic Acid Research* 24(22):4592-4593.
- 35 Yen, H. *et al.* (1997) "The tomato high pigment (*hp*) locus maps to chromosome 2 and influences plastome copy number and fruit quality" *Theoretical and Applied Genetics* 95:1069-1079.
- 40 Yen, H. *et al.* (1995) "The tomato Never-ripe locus regulates ethylene-inducible gene expression and is linked to a homologue of the *Arabidopsis* ETR1 gene" *Plant Physiology* 107:1343-1353.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. ER 716411626 US

INVENTOR(S)				
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)		
Harry J. Denise	Klee Tieman	Gainesville, Florida Gainesville, Florida		19587 U.S. PTO 60/558504
Additional inventors are being named on the _____ separately numbered sheets attached hereto				
TITLE OF THE INVENTION (500 characters max)				
Materials and Methods for Synthesis of a Flavor and Aroma Volatile in Plants				
Direct all correspondence to: CORRESPONDENCE ADDRESS				
<input checked="" type="checkbox"/> Customer Number <div style="border: 1px solid black; padding: 2px; display: inline-block;">23557</div>				
OR				
<input type="checkbox"/> Firm or <input type="checkbox"/> Individual Name				
Address				
Address				
City	State	ZIP		
Country	Telephone	Fax		
ENCLOSED APPLICATION PARTS (check all that apply)				
<input checked="" type="checkbox"/> Specification Number of Pages <u>44</u>		<input type="checkbox"/> CD(s), Number _____		
<input checked="" type="checkbox"/> Drawing(s) Number of Sheets <u>23</u>		<input checked="" type="checkbox"/> Other (specify) <u>Sequence Listing (12 pages)</u>		
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76				
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT				
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.		FILING FEE AMOUNT (\$) <div style="border: 1px solid black; padding: 5px; display: inline-block;">\$80.00</div>		
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees.				
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: <u>19-0065</u>				
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.				
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.				
<input type="checkbox"/> No.				
<input checked="" type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: <u>National Science Foundation</u> <u>Grant Number 7223426-12</u>				

Respectfully submitted,

SIGNATURE

TYPED or PRINTED NAME Doran R. Pace

TELEPHONE (352) 375-8100

Date March 31, 2004

REGISTRATION NO. 38,261

(if appropriate)~

Docket Number: UF-386CP

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Provisional Application
Docket No. UF-386CP

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Harry J. Klee, Denise Tieman
Docket No. : UF-386CP
For : Materials and Methods for Synthesis of a Flavor and Aroma Volatile in Plants

Mail Stop PROVISIONAL APPLICATION
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313

CERTIFICATE OF MAILING BY EXPRESS MAIL (37 CFR §1.10)

Express Mail No.: **ER716411626US** Date of Deposit: March 31, 2004

I hereby certify that the attached Provisional Application and Cover Sheet therefor, with copies as required for authorization for use of Deposit Account No. 19-0065, are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR §1.10 on the date indicated above and are addressed to: Mail Stop PROVISIONAL APPLICATION, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313.

Betty Audette
Name of person mailing paper

Betty Audette
Signature

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2004/032599

International filing date: 01 October 2004 (01.10.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/558,504
Filing date: 31 March 2004 (31.03.2004)

Date of receipt at the International Bureau: 07 March 2006 (07.03.2006)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse